Degree Lower Bounds of Tower-Type for Approximating Formulas with Parity Quantifiers

ALBERT ATSERIAS, Universitat Politècnica de Catalunya
ANUJ DAWAR, University of Cambridge

Kolaitis and Kopparty have shown that for any first-order formula with parity quantifiers over the language of graphs, there is a family of multivariate polynomials of constant-degree that agree with the formula on all but a $2^{-\Omega(n)}$ -fraction of the graphs with n vertices. The proof bounds the degree of the polynomials by a tower of exponentials whose height is the nesting depth of parity quantifiers in the formula. We show that this tower-type dependence is necessary. We build a family of formulas of depth q whose approximating polynomials must have degree bounded from below by a tower of exponentials of height proportional to q . Our proof has two main parts. First, we adapt and extend the results by Kolaitis and Kopparty that describe the joint distribution of the parities of the numbers of copies of small subgraphs in a random graph to the setting of graphs of growing size. Second, we analyze a variant of Karp's graph canonical labeling algorithm and exploit its massive parallelism to get a formula of low depth that defines an almost canonical pre-order on a random graph.

Categories and Subject Descriptors: F.4.1 [Mathematical Logic and Formal Languages]: Mathematical Logic; F.1.3 [Computation by Abstract Devices]: Complexity Measures and Classes; G.2.2 [Discrete Mathematics]: Graph Theory; G.3 [Mathematics of Computing]: Probability and Statistics

General Terms: Algorithms, Theory

Additional Key Words and Phrases: Canonical labeling algorithm, convergence laws, gowers uniformity norm, logic, parity quantifiers, random graphs

ACM Reference Format:

Albert Atserias and Anuj Dawar. 2014. Degree lower bounds of tower-type for approximating formulas with parity quantifiers. *ACM Trans. Comput. Logic* 15, 1, Article 6 (February 2014), 24 pages.
DOI: <http://dx.doi.org/10.1145/2559948>

1. INTRODUCTION

Since the 0-1 law for first-order logic was established [Fagin 1976; Glebskii et al. 1969], there has been much interest in exploring the asymptotic properties of definable classes of graphs. Many extensions of first-order logic have been shown to have a 0-1 law (see for instance Kolaitis and Vardi [1992], Dawar and Grädel [2010]) and in many other cases, weaker forms of convergence have been established (see Compton [1989]). A recent, remarkable result in this vein is that of Kolaitis and Kopparty [2009a] who study $\text{FO}[\oplus]$, the extension of first-order logic with *parity quantifiers*. They show that

A preliminary version of this article appeared in *Proceedings of the 39th International Colloquium on Automata, Languages and Programming (ICALP)*.

A. Atserias was partially supported by CICYT TIN2010-20967-C04-04 (TASSAT). A. Dawar was partially supported by EPSRC grant EP/H026835.

Authors' addresses: Albert Atserias (corresponding author), Universitat Politècnica de Catalunya, Departament de Llenguatges i Sistemes Informàtics c Jordi Girona Salgado 1-3, Edif. Omega-S122 08034 Barcelona, Spain; Anuj Dawar, University of Cambridge Computer Laboratory, 15 J.J. Thomson Avenue, Cambridge CB3 0FD, Great Britain; corresponding author's email: atserias@lsi.upc.edu.

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the Owner/Author.

2014 Copyright is held by the author/owner(s)

1529-3785/2014/02-ART6

DOI: <http://dx.doi.org/10.1145/2559948>

for every constant edge-probability p and for every $\text{FO}[\oplus]$ -sentence ϕ , there are two explicitly computable rational numbers a_0, a_1 such that, for $i \in \{0, 1\}$, as n approaches infinity, the probability that the random graph $G(2n + i; p)$ satisfies ϕ approaches a_i . In other words, ϕ has an asymptotic probability a_0 on the sequence of graphs of even cardinality and a_1 on the sequence of those of odd cardinality. The proof of this result brings entirely new methods to the analysis of the asymptotic behavior of logics on graphs, based on discrete analysis and polynomials over finite fields. In particular, it ties this to the study of approximations of circuits by low-degree polynomials, as we explain next.

The 0-1 law for first-order logic, in its general form, is a quantifier-elimination result. It states that for any first-order formula ϕ , there is a quantifier-free formula θ such that ϕ is equivalent to θ almost surely. To be precise, ϕ and θ agree on a fraction $1 - 2^{-\epsilon n}$ of the graphs on n vertices. We can say that any first-order formula is well approximated by a quantifier-free formula. This is similar to the phenomenon of depth-reduction for circuits which has a long history in computational complexity theory. For instance, Allender showed that AC^0 -circuits have equivalent TC^0 -circuits of depth 3 and quasi-polynomial size [Allender 1989]. The result of Beigel and Tarui that general ACC^0 -circuits have equivalent depth-2 circuits of quasi-polynomial size with a symmetric gate at the root [Beigel and Tarui 1994] has been exploited to remarkable effect recently in the work of Williams [2011]. In the context of approximation, one of the best known examples is the Razborov-Smolensky approximation of $\text{AC}^0[\oplus]$ -circuits by multivariate polynomials over $\mathbb{Z}_2$ of polylogarithmic degree [Razborov 1987; Smolensky 1987]. The method yields an approximation that agrees on a fraction $1 - 2^{-(\log n)^c}$ of the inputs.

The Kolaitis-Kopparty result previously mentioned is proved by a depth-reduction argument of a similar kind that exploits the higher degree of symmetry that $\text{FO}[\oplus]$ -formulas have over $\text{AC}^0[\oplus]$ -circuits. They prove that every $\text{FO}[\oplus]$ -formula ϕ is well-approximated by a formula which is a Boolean combination of quantifier-free formulas and of polynomials over $\mathbb{Z}_2$ of special form, that we call *$\text{FO}[\oplus]$ -polynomials*, and the degree of these polynomials depends only on ϕ . These polynomials have as variables X_{uv} for every potential edge $\{u, v\}$ over the vertex-set $\{1, \dots, n\}$. For example, the $\text{FO}[\oplus]$ -polynomial that gives the parity of the number of triples that extend the vertex u to a triangle is

$$\sum_{\substack{v: \\ v \neq u}} \sum_{\substack{w: \\ w \neq u \\ w \neq v}} X_{uv} X_{vw} X_{wu}. \quad (1)$$

At the heart of the argument is the analysis of the bias of certain low-degree polynomials of this type on uniformly random inputs. This understanding is then used to carry over a quantifier-elimination argument that eliminates one parity quantifier or one first-order quantifier at a time. Relevant to our work is the fact that, intriguingly, the elimination of each parity quantifier in this argument incurs an exponential loss in the degree. The final outcome is that the degree d of the approximating polynomials is bounded from below by a function of tower-type on the number q of parity quantifiers that were eliminated, that is,

$$d \geq 2^{2^q}, \quad (2)$$

where the height of the tower is at least q . At first sight, the source of this inefficiency in the proof appears technical, and it might be tempting to think that a different method could perhaps avoid it altogether.

In this article, we prove that the non-elementary dependence stated in Equation (2) cannot be avoided. To be precise, we construct an explicit family of $\text{FO}[\oplus]$ -formulas ϕ_q of quantifier rank q and prove that they cannot be approximated by a Boolean

combination of quantifier-free formulas and polynomials of degree bounded by an elementary function of q . Specifically, we prove the following.

THEOREM 1.1. *There exists a constant $c > 0$ such that for every large enough integer q , every $\epsilon > 0$, and every large enough integer n , there exists an $\text{FO}[\oplus]$ -formula $\phi(u, v, w)$ of quantifier rank q such that for every Boolean combination p of quantifier-free formulas and $\text{FO}[\oplus]$ -polynomials of degree bounded by a tower of exponentials of height at most q/c , the formulas ϕ and p must disagree on a fraction $1 - \epsilon$ of all graphs with n vertices.*

By an $\text{FO}[\oplus]$ -polynomial we mean a formula that has a direct translation to a bounded-degree polynomial over $\mathbb{Z}_2$: a sequence of parity quantifiers followed by a conjunction of atomic facts. Since both ϕ and p have free variables, when we say that ϕ and p disagree on a graph G , we mean that the set of tuples of vertices of G satisfying ϕ is not the same as the set of tuples satisfying p .

Theorem 1.1 should be contrasted with the 0-1 law for first-order logic. In that case, the approximating formula is quantifier-free, and such formulas translate into polynomials of degree at most polynomial in the number of free variables.

Proof Outline and Techniques. Our proof relies on two technical ingredients. On one hand, we analyze a canonical labeling algorithm for graphs due to Karp [1979] (see [Hella et al. 1996] for another view on the logical definability of Karp's canonical labeling). We exploit its massive parallelism to build an $\text{FO}[\oplus]$ -formula $\psi(u, v)$ of depth $O(\log^* n)$ that works on graphs with n vertices. The formula is designed in such a way that, on almost every graph, it defines a linear pre-order of width at most two on the set of vertices of the graph. The second ingredient is a refined analysis of one of the key tools from the Kolaitis-Kopparty paper. Using and extending their techniques for estimating the frequencies mod 2 of subgraph copies, we show that for every $\text{FO}[\oplus]$ -polynomial $p(u, v, w)$ of degree $\log \log \log n$ and for the random graph $G(n, 1/2)$, with high probability, there exists a triple of distinct vertices (a, b, c) such that p cannot distinguish it from any of its permutations.

From these two ingredients, the lower bound follows by taking the formula $\phi(u, v, w) := \psi(u, v) \wedge \psi(v, w)$. On one hand this formula distinguishes at least one permutation of the vertices (a, b, c) from some other. This is because by linearity of the pre-order, the classes they lie in must be comparable, but by the width-2 condition on the pre-order, not all three vertices can sit in the same class. On the other hand, if $\phi'(u, v, w)$ is any Boolean combination of quantifier-free formulas and $\text{FO}[\oplus]$ -polynomials of degree $\log \log \log n$, we could choose (a, b, c) in such a way that ϕ' is not able to distinguish any permutation of (a, b, c) from the others. We conclude that ϕ' cannot approximate ϕ , and since the quantifier rank of ϕ is still $O(\log^* n)$, the tower-type lower bound follows. We provide more details in the body of the article.

Section 2 introduces some essential notation. Then, in Section 3, we show that for any Boolean combination of polynomials of low degree, on a sufficiently large random graph, there is some tuple of elements which is not distinguished from any of its permutations. Section 4 contains the construction of the formula $\psi(u, v)$ that defines a linear pre-order of width at most two on almost all graphs on n vertices. Finally, Section 5 pulls these ingredients together to establish our result. The construction in Section 3 relies heavily on elements from Kolaitis and Kopparty [2009a, 2009b]. Similarly, Section 4 borrows from the analysis in Karp [1979]. The reader wishing to follow all details of the proof may wish to have copies of these three papers at hand.

2. PRELIMINARIES

We use $[n]$ to denote the set $\{1, \dots, n\}$. We identify the nodes of a complete rooted binary tree with the binary strings that start with the symbol 1: the root is 1, the left

child of t is $t0$, and the right child of t is $t1$. The *level order* of a complete binary tree is 1, 10, 11, 100, 101, 110, 111, 1000, 1001, $\dots$, that is, the nodes are ordered first by length, and within each length, they are ordered lexicographically. Note that if the strings are interpreted as numbers written in binary, this is the usual order of the natural numbers. For a natural number $n \geq 1$, we write $\text{bin}_2(n)$ for its unique binary encoding with a leading one.

Let G and H be graphs. We write $V(G)$ and $E(G)$ to denote the vertices and edges of G , respectively, and similarly for H . A homomorphism from G to H is a mapping $h : V(G) \rightarrow V(H)$ that maps edges to edges; that is, such that if $\{u, v\} \in E(G)$, then $\{h(u), h(v)\} \in E(H)$. Let $\text{Hom}(G, H)$ denote the collection of all homomorphisms from G to H .

The collection of $\text{FO}[\oplus]$ -formulas over the language of graphs is the smallest class of formulas that contains the atomic formulas $E(x, y)$ and the equalities $x = y$, and is closed under negation, conjunction and disjunction, universal and existential quantification, and parity quantification; that is, quantification of the form $\oplus x \phi(x)$. The meaning of $\oplus x \phi(x)$ is that there is an odd number of vertices x that satisfy $\phi(x)$. For a tuple $\mathbf{a} = (a_1, \dots, a_k)$ and a permutation $\pi \in S_k$, we write $\mathbf{a} \circ \pi$ for the tuple $(a_{\pi(1)}, \dots, a_{\pi(k)})$. If $p(x_1, \dots, x_k)$ is a formula with free variables $x_1, \dots, x_k$, and $y_1, \dots, y_k$ are variables or constants, we write $p(y_1, \dots, y_k)$ for the result of replacing each occurrence of x_i by y_i . This applies also to the case where $y_1, \dots, y_k$ is a permutation of $x_1, \dots, x_k$.

An *atomic type* on the variables $x_1, \dots, x_k$ over the language of graphs is a consistent collection of atomic formulas $E(x_i, x_j)$ or $x_i = x_j$ and negated atomic formulas $\neg E(x_i, x_j)$ or $x_i \neq x_j$ that is maximal with respect to set-inclusion. A *positive atomic type* is the subset of an atomic type containing all its positive atomic formulas. We say that a type is *injective* if it contains the formula $x_i \neq x_j$ whenever $i \neq j$. An *equality type* is the subset of an atomic type containing all its equalities $x_i = x_j$ and inequalities $x_i \neq x_j$. For a graph G and a tuple $\mathbf{a} = (a_1, \dots, a_k) \in V(G)^k$, the atomic type of $\mathbf{a}$ in G is the unique atomic type that is made true in G by the assignment $x_i \mapsto a_i$. The atomic type of a_k over $(a_1, \dots, a_{k-1})$ in G is the subset of the atomic type of $\mathbf{a}$ in G containing all formulas that involve the variable x_k .

3. FOOLING POLYNOMIALS OF LOW DEGREE

In this section, we aim to establish that for any $\text{FO}[\oplus]$ -formula which is a Boolean combination of polynomials of low degree (growing as $O(\log \log \log n)$) and a sufficiently large random graph G , with high probability there is a triple of vertices a, b, c such that the formula does not distinguish this triple from any of its permutations. To do this, we first establish a normal form for such $\text{FO}[\oplus]$ -formulas which will permit an analysis of their asymptotic behavior. The normal form is established in Section 3.3, and the analysis in Section 3.4.

Once we have the normal form, the proof strategy is roughly as follows. Fix a formula in normal form $p(x, y, z)$. For every fixed $a, b, c \in [n]$, let $Y(a, b, c)$ be the event that p cannot distinguish any two permutations of a, b, c . Ideally we would like to show that the event $Y(a, b, c)$ has nonnegligible probability of happening, and that if $a', b', c' \in [n]$ is a triple disjoint from a, b, c , then the events $Y(a, b, c)$ and $Y(a', b', c')$ are almost independent. If we were able to do this, the result would follow from an application of Chebyshev's inequality. Unfortunately it is not quite true that $Y(a, b, c)$ and $Y(a', b', c')$ are almost independent in general, so we need to take a detour. The detailed argument is given in Section 3.5

3.1. Formulas and Polynomials

In this section, we define the formulas to which our result applies. In short, they are Boolean combinations of $\text{FO}[\oplus]$ -polynomials. An $\text{FO}[\oplus]$ -polynomial is a formula of

FO[$\oplus$] consisting of a sequence of parity quantifiers followed by a conjunction of atomic formulas and negated equalities. In its general form, an FO[$\oplus$]-polynomial p with free variables $u_1, \dots, u_k$ is a formula of the form

$$\oplus u_{k+1} \cdots \oplus u_m \left(\bigwedge_{i \neq j} u_i \neq u_j \wedge \bigwedge_{\ell=1}^d E(u_{i_\ell}, u_{j_\ell}) \right), \quad (3)$$

where i and j range over $[m]$ in the first conjunction, and $i_1, \dots, i_d, j_1, \dots, j_d$ are indices in $[m]$. The number d of atomic facts in the conjunction is called the *degree* of p . The number m of distinct variables is the *order* of p .

To Formula (3), we can associate the graph H on the vertices $u_1, \dots, u_m$ that has an edge between u_{i_ℓ} and u_{j_ℓ} for each $\ell \in [d]$, and the first k vertices $u_1, \dots, u_k$ distinguished from the rest. We use the notation $H(u_1, \dots, u_k)$ for such kind of graphs, and $\oplus H(u_1, \dots, u_k)$ to denote the FO[$\oplus$]-polynomial corresponding to this graph. Note that the formula expresses the parity of the number of extensions of $u_1, \dots, u_k$ to a copy of H . Note also that the degree of $\oplus H(u_1, \dots, u_k)$ is the number of edges of H .

Example 3.1. If H is a triangle containing vertex u , then $\oplus H(u)$ is the formula that expresses the parity of the number of extensions of u to a triangle. Formally, $\oplus H(u)$ is the formula

$$\oplus v \oplus w (u \neq v \wedge u \neq w \wedge v \neq w \wedge E(u, v) \wedge E(v, w) \wedge E(w, u)).$$

Note that over undirected graphs, this formula is always false. This is because for every triangle containing u , there are two assignments to the variables v and w which witness H . Thus, the total number of satisfying assignments is twice the number of triangles containing u and is therefore always even. In general, if H has an even number of automorphisms that fix $u_1, \dots, u_k$, then $\oplus H(u_1, \dots, u_k)$ will always be false, while for graphs with an odd number of automorphisms, we get nontrivial formulas.

Remark 3.2. The observation at the end of Example 3.1 has one important consequence: if $H(u_1, \dots, u_k)$ has r isolated vertices outside $\{u_1, \dots, u_k\}$ and $r \geq 2$, then $\oplus H(u_1, \dots, u_k)$ is always false, because then the number of automorphisms of H that fix $u_1, \dots, u_k$ is a multiple of $r!$, which is even when $r \geq 2$. In particular, this means that in the general form of FO[$\oplus$]-polynomials displayed in Equation (3), we can always assume that

$$m \leq 2d + 1.$$

This will be important because, by assuming it, lower bounds on the order imply lower bounds on the degree.

Remark 3.3. On graphs, an atomic formula of the form $E(u_1, u_1)$ is just false, while an atomic formula of the form $E(u_1, u_2)$ with distinct variables u_1 and u_2 is equivalent to $u_1 \neq u_2 \wedge E(u_1, u_2)$, which is an FO[$\oplus$]-polynomial of order two and degree one. Similarly, an atomic formula of the form $u_1 = u_1$ is just true, while an atomic formula of the form $u_1 = u_2$ with distinct variables u_1 and u_2 is equivalent to the negation of the FO[$\oplus$]-polynomial $u_1 \neq u_2$; this has order two and degree zero. This means that every quantifier-free formula is equivalent to a Boolean combination of FO[$\oplus$]-polynomials of order two and degree at most one.

Remark 3.4. There is a precise sense in which FO[$\oplus$]-polynomials correspond to polynomials over the Boolean edge-variables X_{uv} . For example, the formula from

Example 3.1 corresponds to the family of degree-3 polynomials

$$\sum_{\substack{v \in [n] \\ v \neq u}} \sum_{\substack{w \in [n] \\ w \neq u \\ w \neq v}} X_{uv} X_{vw} X_{wu}.$$

as u ranges over $[n]$.

3.2. Independence and Plan of Action

The formulas we are interested in are Boolean combinations of polynomials. Let $p(x, y, z)$ be such a formula and $Y(a, b, c)$ be the event that p cannot distinguish any two permutations of a, b, c . The main obstacle to carrying out the argument sketched at the beginning of this section is that it is not true, in general, that the events $Y(a, b, c)$ and $Y(a', b', c')$ are almost independent, even if a, b, c, a', b', c' are all different. The reason is that the formula $p(x, y, z)$ may include statements about the graph G which do not involve the free variables. These are true or false independently of the choice of a, b, c or a', b', c' and thus create correlations between $Y(a, b, c)$ and $Y(a', b', c')$.

It is illustrative to give an example of this.

Example 3.5. Let $p(x, y)$ be the formula that is the conjunction of the following: (1) $\oplus z E(x, z)$ (x has odd degree); (2) $\neg \oplus z E(y, z)$ (y has even degree); and (3) $\oplus H$ for some fixed nontrivial rigid graph H (x and y do not appear free in this). This is a Boolean combination of $\text{FO}[\oplus]$ -polynomials of degree bounded by the number of edges of H .

Note that if $p(a, b)$ holds, then $p(b, a)$ must fail. Therefore, the probability that $p(a, b) \not\leftrightarrow p(b, a)$ holds is approximately $2 \cdot \frac{1}{8}$, since each of the three conditions in $p(a, b)$ holds with probability approximately $\frac{1}{2}$ almost independently, and similarly for $p(b, a)$. On the other hand, the probability that both $p(a, b) \not\leftrightarrow p(b, a)$ and $p(a', b') \not\leftrightarrow p(b', a')$ hold simultaneously is approximately $4 \cdot \frac{1}{32}$. This is because in each of the four cases in which both hold, condition (3) either holds for both a, b and a', b' or for neither (since x and y do not appear). We are left with five conditions that hold with probability approximately $\frac{1}{2}$ almost independently. Since $4 \cdot \frac{1}{32}$ is not ϵ -close to $(2 \cdot \frac{1}{8})^2$, this shows that $Y(a, b)$ and $Y(a', b')$ are not almost independent.

The example just sketched suggests that we factor out the condition that does not depend on neither x nor y from $p(x, y) \leftrightarrow p(y, x)$, since this is the cause for the statistical dependence between $Y(a, b)$ and $Y(a', b')$. However, while such an argument can be made to work in the preceding example, it is not clear what such a factoring would entail when p contains disjunctions.

The key observation at this point is that the *full type* of (x, y) in terms of its atomic type (the pattern of connections and equalities among x and y) and the truth values of its $\oplus H$'s as H ranges over all small graphs that contain x and y as vertices is enough to determine the truth value of $p(x, y)$. Thus, if we were able to find a full type implying $p(x, y)$ that is symmetric in x and y , we would have reduced the case of general $p(x, y)$ to the case of a $p(x, y)$ that consists of a single term and eliminated the need to consider disjunctions. The argument that we use is a bit more delicate than this, but this is the main idea.

3.3. Normal Forms

In this section, we introduce some definitions and discuss two different types of *normal forms* for Boolean combinations of $\text{FO}[\oplus]$ -polynomials.

An *I-labeled graph* is a graph with some vertices labeled by elements of I in such a way that, for every $i \in I$, there is exactly one vertex labeled i , and the set of labeled vertices induces an independent set. The set of labeled vertices of an I -labeled graph

H is denoted by $\mathcal{L}(H)$. The vertex labeled by $i \in I$ is denoted by $H(i)$. An I -labeled graph H is *label-connected* if $H \setminus \mathcal{L}(H)$ is connected. Let Conn_I^t be the set of all I -labeled label-connected graphs with at most t unlabeled vertices. We say that H *depends on* label $i \in I$ if $H(i)$ is not an isolated node. We say that H is *label-dependent* if it depends on all its labels. Let $\text{Conn}_I^{*,t}$ be the subset of all labeled graphs in Conn_I^t that are label-dependent.

A k -labeled graph is a $[k]$ -labeled graph. A $\leq k$ -labeled graph is an I -labeled graph for some $I \subseteq [k]$. Let H be a $\leq k$ -labeled graph with labels $I \subseteq [k]$. A *homomorphism* from H to a pair $(G, \mathbf{a})$, where G is a graph and $\mathbf{a} = (a_1, \dots, a_k)$ is a tuple in $V(G)^k$, is a homomorphism $\chi \in \text{Hom}(H, G)$ such that $\chi(H(i)) = a_i$ for each $i \in I$. A homomorphism χ from H to $(G, \mathbf{a})$ is *injective* if for any distinct $a, b \in V(H)$ such that $\{a, b\} \not\subseteq \mathcal{L}(H)$, we have $\chi(a) \neq \chi(b)$. Write $\oplus H(G, \mathbf{a})$ for the parity of the number of injective homomorphisms from H to $(G, \mathbf{a})$. We usually omit G and write $\oplus H(\mathbf{a})$. When H is a k -labeled graph (i.e., $I = [k]$), the notation for this in Kolaitis and Kopparty [2009a] is $[H]_2(G, \mathbf{a})$.

We call the number of vertices in H the *order* of H and the number of edges in H the *degree* of H . These are the same as the order and degree, respectively, of the polynomial $\oplus H(\mathbf{x})$. Note that if H has order c , then it has at most c unlabeled vertices.

A *KK-normal form* of order m with free-variables $\mathbf{x} = (x_1, \dots, x_k)$ is a Boolean combination of the atomic types on the variables $\mathbf{x}$ and formulas $\oplus H(\mathbf{x})$'s as H ranges over the k -labeled label-connected graphs of order m with labeled vertices $\mathbf{x}$. A *regular normal form* of order m with free-variables $\mathbf{x}$ is a Boolean combination of the atomic types on the variables $\mathbf{x}$ and the $\oplus H(\mathbf{x})$'s as H ranges over the $\leq k$ -labeled label-connected, label-dependent graphs of order m with labeled vertices within $\mathbf{x}$.

Example 3.6. Let $\phi(x, y)$ be the formula

$$\oplus z (E(x, z)) \wedge \neg \oplus z (E(y, z)),$$

saying that x has odd degree and y has even degree. This is a regular normal form. On the other hand, it is *not* a KK-normal form, because the formula $\oplus z (E(x, z))$ cannot be put in the form $\oplus H(x, y)$ for any 2-labeled graph H . However, as we will see, it is not hard to transform $\phi(x, y)$ into an equivalent KK-normal form.

Example 3.7. Let $p(x, y)$ be the formula

$$(x \neq y \wedge E(x, y) \wedge \neg \oplus H_1(x, y) \wedge \oplus H_2(x, y)) \vee (x \neq y \wedge \neg E(x, y) \wedge \oplus H_1(x, y) \wedge \neg \oplus H_2(x, y)),$$

where H_1 is the 2-labeled label-connected graph that has three vertices x, y , and z and a single edge between x and z , and H_2 is the 2-labeled label-connected graph that has three vertices x, y , and z and a single edge between y and z . This is a KK-normal form. On the other hand, it is not a regular normal form, because H_1 and H_2 are not label-dependent. However, as we will see, it is not hard to transform $\phi(x, y)$ into a regular normal form.

These two examples are actually logically equivalent, and it is a general fact that Boolean combinations of $\text{FO}[\oplus]$ -polynomials, KK-normal forms, and regular normal forms of the same order have the same expressive power.

LEMMA 3.8. *Let $k \geq 0$ and $m \geq k$ be integers and let $\phi(x_1, \dots, x_k)$ be an $\text{FO}[\oplus]$ -formula. The following are equivalent.*

- (1) ϕ is equivalent to a Boolean combination of $\text{FO}[\oplus]$ -polynomials of order at most m .
- (2) ϕ is equivalent to a KK-normal form of order at most m .
- (3) ϕ is equivalent to a regular normal form of order at most m .

PROOF. We may assume that $m \geq 2$, since $\text{FO}[\oplus]$ -polynomials and normal forms of order one or less are trivial.

(1) $\Rightarrow$ (2). We show how to transform an FO[$\oplus$]-polynomial p of order at most m into an equivalent Boolean combination of atomic types and formulas of the form $\oplus H$, where H is a k -labeled label-connected graph of order at most m . We do this in two steps: in the first step, we do it with k -labeled graphs that are not necessarily label-connected, and in the second step, we ensure the connectivity condition.

Let us say p has the form

$$p(y_1, \dots, y_t) = \oplus y_{t+1} \cdots \oplus y_s \left(\bigwedge_{i \neq j} y_i \neq y_j \wedge \bigwedge_{\ell=1}^d E(y_{i_\ell}, y_{j_\ell}) \right), \quad (4)$$

where the indices i and j range over $[s]$ in the first conjunct, i_ℓ and j_ℓ are also indices in $[s]$ for each $\ell \in [d]$, the free variables $y_1, \dots, y_t$ are among $x_1, \dots, x_k$, and the bound variables $y_{t+1}, \dots, y_s$ are disjoint from $x_1, \dots, x_k$. After renaming the variables, we may assume that $(y_1, \dots, y_t) = (x_1, \dots, x_t)$. Let H be the k -labeled graph of order s with labeled vertices $x_1, \dots, x_k$, non-labeled vertices $y_{t+1}, \dots, y_s$, and edges $\{y_{i_\ell}, y_{j_\ell}\}$ as ℓ ranges over $[d]$. We claim that for every graph G and every $\mathbf{a} = (a_1, \dots, a_k) \in V_G^k$, we have

$$p(\mathbf{a}) = \sum_{f \in \mathcal{F}} \sum_{\sigma \in S^f} \sigma(\mathbf{a}) \cdot \oplus H^f(\mathbf{a}) \pmod{2}, \quad (5)$$

where the following hold.

- $\mathcal{F}$ is the set of partial mappings $f : [s] \setminus [t] \rightarrow [k] \setminus [t]$ that are injective on their domain $\text{Dom}(f)$, including the empty map.
- For every $f \in \mathcal{F}$, the set S^f consists of all atomic types on the variables $x_1, \dots, x_k$ that contain the positive atomic type of each y_j with $j \in \text{Im}(f)$ over $x_1, \dots, x_t$ in H , plus the inequalities $x_i \neq x_j$ for every pair $i, j \in [t]$ with $i \neq j$, every pair $i, j \in [k]$ with $i \in [t]$ and $j \in \text{Im}(f)$, and every pair $i, j \in \text{Im}(f)$ with $i \neq j$.
- For every $f \in \mathcal{F}$, the graph H^f is the k -labeled graph that is obtained from H by deleting all edges between some y_i with $i \in \text{Dom}(f)$ and $x_1, \dots, x_t$, and by identifying y_i with $x_{f(i)}$ for every $i \in \text{Dom}(f)$.

To see that Eq. (5) holds, note that the term $\sigma(\mathbf{a}) \cdot \oplus H^f(\mathbf{a})$ counts the parity of the number of injective homomorphisms from H to G that map x_i to a_i for every $i \in [t]$, and y_i to $a_{f(i)}$ for every $i \in \text{Dom}(f)$, subject to the condition that $\mathbf{a}$ satisfies the rest of atomic relations specified in σ . Since for each $f \in \mathcal{F}$ all allowed possibilities for the rest of atomic relations are included, the resulting count is precisely $p(\mathbf{a})$. Now note that $\mathcal{F}$ and S^f are finite for every $f \in \mathcal{F}$, and therefore Expression (5) is equivalent to a Boolean combination of atomic types and $\oplus H$'s, as required.

It remains to see how to ensure that the H 's are label-connected while preserving the bound on the order. Conveniently, this was done in Lemma 5.6 from Kolaitis and Kopparty [2009a], and we refer the reader to it.

(2) $\Rightarrow$ (3). We need to show how to transform a formula $\oplus H$, where H is a k -labeled label-connected graph of order at most m into an equivalent Boolean combination of quantifier-free formulas of the form $\oplus F$, where F is a $\leq k$ -labeled label-connected, label-dependent graph of order at most m . The transformation is done in two steps. In the first step, we reduce the number of isolated labeled vertices in H or the number of non-labeled vertices of H at the expense of using possibly label-disconnected graphs. In the second step, we get rid of the label-disconnected graphs. Indeed, the second step, is as in the proof of Lemma 5.6 in Kolaitis and Kopparty [2009a], so we need only take care of the first step.

If H is already label-dependent, there is nothing to do. Otherwise let $x_1, \dots, x_k$ be the set of labeled vertices of H , assume x_k is isolated in H , and let $x_{k+1}, \dots, x_s$ be the set of

non-labeled vertices of H . Then, for every graph G and for every $\mathbf{a} = (a_1, \dots, a_k) \in V_G^k$, we claim that

$$\oplus H(\mathbf{a}) \equiv \oplus H^{-x_k}(\mathbf{a}) + \sum_{j \in [s] \setminus [k]} \sum_{\sigma \in S^j} \sigma(\mathbf{a}) \cdot \oplus H^{x_j=x_k}(\mathbf{a}) \pmod{2}, \quad (6)$$

where the following hold.

- H^{-x_k} is the $\leq k$ -labeled graph that results from deleting x_k in H .
- S^j is the set of atomic types on the variables $x_1, \dots, x_k$ that contain the positive atomic type of x_j over $x_1, \dots, x_{k-1}$ in H , plus the inequalities $x_j \neq x_i$ for every $i \in [k-1]$.
- $H^{x_j=x_k}$ is the k -labeled graph that results from deleting all edges from x_j to a labeled vertex and identifying x_j and x_k in H .

To see why Eq. (6) holds, note two facts. First, the sum over $j \in [s] \setminus [k]$ counts the parity of the number of injective homomorphisms from H^{-x_k} into $(G, \mathbf{a})$ that have a_k as the image of some non-labeled vertex. Second, since the term $\oplus H^{-x_k}(\mathbf{a})$ counts the parity of all injective homomorphisms from H^{-x_k} into $(G, \mathbf{a})$, this means that each injective homomorphism from H^{-x_k} into $(G, \mathbf{a})$ that has a_k as the image of some non-labeled vertex is counted exactly twice and cancels. What is left is the parity of the number of injective homomorphisms from H^{-x_k} into $(G, \mathbf{a})$ that does not have a_k as the image of some non-labeled vertex. This is precisely $\oplus H(\mathbf{a})$, because x_k is isolated in H , and therefore the only constraint it puts on the injective homomorphisms from H into $(G, \mathbf{a})$ is that non-labeled vertices are not mapped to a_k .

To conclude, note that $[s] \setminus [k]$ and S^j are finite for every $j \in [s] \setminus [k]$, and therefore (6) is a Boolean combination of atomic types and $\oplus H$'s as required.

(3) $\Rightarrow$ (1). First, each atomic type is a quantifier-free formula, and hence a Boolean combination of FO[$\oplus$]-polynomials of order at most two and degree at most one by Remark 3.3. Here we use the assumption made at the beginning that $m \geq 2$.

Second, let H be a $\leq k$ -labeled graph of order at most m . Let $y_1, \dots, y_t$ be its set of labeled vertices, which is a subset of $x_1, \dots, x_k$, and let $y_{t+1}, \dots, y_s$ be its set of unlabeled vertices, which is a set disjoint from $x_1, \dots, x_k$. Then

$$\oplus H(\mathbf{x}) \equiv \sum_{\sigma \in S} \sigma(\mathbf{x}) \cdot \oplus H^\sigma(\mathbf{x}) \pmod{2}, \quad (7)$$

where the following hold.

- S is the collection of all equality types on $y_1, \dots, y_t$.
- H^σ is the $\leq k$ -labeled graph that results from H by identifying every vertex y_j with $j \in [t]$ with the vertex y_i with smallest index $i \in [t]$ for which the equality $y_i = y_j$ appears in σ , and by deleting duplicated edges. If $i \neq j$, we say that y_i survives and y_j disappears. All vertices y_i with $i \in [s] \setminus [t]$ survive.

For fixed $\sigma \in S$, let A be the set of pairs $(i, j) \in [s]^2$ such that both vertices y_i and y_j survive in H^σ . Finally, let B be the subset of pairs (i, j) in A such that $\{y_i, y_j\}$ is an edge in H^σ . Then

$$\sigma(\mathbf{x}) \cdot \oplus H^\sigma(\mathbf{x}) \equiv \oplus y_{t+1} \cdots \oplus y_s \left(\bigwedge_{(i,j) \in A} y_i \neq y_j \wedge \bigwedge_{(i,j) \in B} E(y_i, y_j) \right).$$

This is precisely an FO[$\oplus$]-polynomial whose free variables are the y_i with $i \in [t]$ that survive in H^σ . Its order is s , which is at most m . Since S is finite, this shows that Eq. (7) is a equivalent to a Boolean combination of FO[$\oplus$]-polynomials of order at most m . $\square$

3.4. Distribution of Frequency Vectors

The frequency vector of order t in a graph G is the $\{0, 1\}$ -vector indexed by the set of all connected graphs with at most t vertices where the component H is $\oplus H(G)$, that is, the parity of the number of occurrences of H in G . Kolaitis and Kopparty [2009a] give an analysis of the distribution of frequency vectors in a random graph $G \sim G(n, 1/2)$, for constant t . Our aim in the present section is to extend this analysis to orders that grow with n and to $\leq k$ -labeled graphs.

Let $\text{Conn}_{\leq k}^t$ be the set of all $\leq k$ -labeled label-connected graphs with at most t unlabeled vertices. Let $\text{Conn}_{\leq k}^{*,t}$ be the subset of $\text{Conn}_{\leq k}^t$ containing all graphs that are label-dependent. Let G be a graph, let $\mathbf{a}$ be a tuple in $V(G)^k$, and let $t \geq 0$ be an integer. Let $\text{freq}_{\leq k, G}^{*,t}(\mathbf{a})$ be the $\{0, 1\}$ -vector indexed by the elements $\text{Conn}_{\leq k}^{*,t}$ that has $\oplus H(\mathbf{a})$ as its component indexed by H . Next we extend the definition of feasible frequency vectors from Kolaitis and Kopparty [2009a] to the setting of $\leq k$ -labeled graphs. In defining $\text{FFreq}^*(\tau, \leq k, t)$, we will restrict our attention to injective atomic types τ . This simplifies matters significantly and is enough for our purposes. If τ is an injective atomic type on $x_1, \dots, x_k$, let $\text{FFreq}^*(\tau, \leq k, t)$ denote the set of all *feasible frequency vectors*. Explicitly, these are all the $\{0, 1\}$ -vectors indexed by $\text{Conn}_{\leq k}^{*,t}$ whose component F belongs to $\text{aut}(F) \cdot \mathbb{Z}_2$. Here $\text{aut}(F)$ denotes the number of automorphisms of F that fix the labels. Let $\text{FFreq}_n^*(\tau, \leq k, t)$ denote the set of $f \in \text{FFreq}^*(\tau, \leq k, t)$ such that $f_{K_1(\emptyset)} = n \bmod 2$, where $K_1(\emptyset)$ is the graph with no labels and exactly one unlabeled vertex.

The next lemma describes the distribution of $\text{freq}_{\leq k, G}^{*,t}(\mathbf{a})$ in a random graph. This is analogous to Theorem 2.4 in Kolaitis and Kopparty [2009a], but see also the statement of Theorem 8.2 in Kolaitis and Kopparty [2009b] (about subgraph *copies*), on which Theorem 2.4 is based. Our statement deviates from theirs in two directions: in generalizing it from a constant number of vertices to a growing number of vertices up to $\log \log \log n$, and from k -labeled graphs to $\leq k$ -labeled graphs.

LEMMA 3.9. *For every $k \geq 0$, there exists $n_0 \geq 0$ such that for every $n \geq n_0$, every injective atomic type τ on k variables, every $c \leq \log \log \log n$, and every k -tuple $\mathbf{a}$ of distinct elements in $[n]$, the distribution of $\text{freq}_{\leq k, G}^{*,c}(\mathbf{a})$ as $G \sim G(n, 1/2 \mid \tau(\mathbf{a}))$ is $2^{-\Omega_k(n/\log n)}$ -close in statistical distance from the uniform distribution over $\text{FFreq}_n^*(\tau, \leq k, c)$.*

Here, the notation Ω_k refers to an unspecified multiplicative constant that depends only on k , and the notation $G(n, 1/2 \mid \tau(\mathbf{a}))$ denotes the uniform distribution over the graphs with vertex-set $[n]$ restricted to those on which the set of vertices in the tuple $\mathbf{a}$ induces the subgraph specified by the atomic type τ .

Before we discuss its proof, it is worth pointing out the differences between the statement of Lemma 3.9 and the statements of Theorem 2.4 in Kolaitis and Kopparty [2009a] and the more general Theorem 8.2 in Kolaitis and Kopparty [2009b]. Our statement here extends them in two directions, and both of these require significant adaptation of the proof.

The first difference concerns the extension which takes us from a bounded number of vertices to a growing number of vertices up to $\log \log \log n$. In order to achieve this, we relax the statistical distance from $2^{-\Omega_k(n)}$ in Kolaitis and Kopparty [2009b] to $2^{-\Omega_k(n/\log n)}$ here. For our purposes, this weaker bound on the distance is not significant. The proof is obtained by adapting the one in Kolaitis and Kopparty [2009b]. For this we need to make an explicit calculation of an ϵ -bound in Lemma 4.7 in Kolaitis and Kopparty [2009b]. It is conceivable that a more careful analysis would yield a better bound still. However, the bound we get is sufficient for our purposes.

The second difference concerns the extension from k -labeled graphs in Theorem 8.2 in Kolaitis and Kopparty [2009b] to $\leq k$ -labeled graphs here. We need this extension

to be able to execute the *factoring* argument sketched in Section 3.2. This modification in the statement introduces the additional restriction of label-dependency (the $*$ in $\text{freq}^{*,t}$). This is required, since the result would not be true without it. Luckily, though, Lemma 3.8 tells us that we will be able to assume label-dependency without loss of generality. In this case, discovering the right assumption is the essential step in the proof. Once the concept is defined, the proof again follows the original one. One final difference between our statement and Theorem 8.2 in Kolaitis and Kopparty [2009b] is that we are stating only the case $s = \ell' = 0$ of their theorem, since any other case is irrelevant for our application. This also has the benefit of shortening the proof somewhat.

PROOF (SKETCH OF LEMMA 3.9). Handling $\leq k$ -labeled graphs instead of k -labeled graphs in the proof of Theorem 8.2 in Kolaitis and Kopparty [2009b] is not problematic until we realize that the sets of copies of F and F' in $(K_n, \mathbf{a})$ need not be disjoint, even if F and F' are non-isomorphic; that is, Proposition 8.1(2) from Kolaitis and Kopparty [2009b] fails if $I = [k]$ and F and F' are $\leq k$ -labeled graphs instead of k -labeled graphs. This happens, for example, if F and F' are $\leq k$ -labeled label-connected graphs that are identical, except that F' has one more labeled vertex than F that is isolated. On the other hand, if F and F' are non-isomorphic and depend on all its labels, then it can be seen that the sets of copies are disjoint. This is enough to carry over the argument in Kolaitis and Kopparty [2009a].

However, in order to allow a growing c , we need the following lemma that makes the ϵ -bound explicit in the conclusion of Lemma 4.7 in Kolaitis and Kopparty [2009b]. For the definition of the μ -Gowers norm $\|g\|_{U^d, \mu}$, see Section 4.1 in Kolaitis and Kopparty [2009b].

LEMMA 3.10. *Let $g : \mathbb{Z}_2^d \rightarrow \{-1, 1\}$ be given by $g(\mathbf{y}) = (-1)^{\prod_{i=1}^d y_i}$. Let μ be the uniform distribution on $\mathbb{Z}_2^d$. Then $\|g\|_{U^d, \mu} < 1 - \epsilon$, where $\epsilon > 1 - \exp(-2^{-d^2-2d+1})$.*

PROOF. Let $p_0 = \mu^{(d)}(\mathbf{0}, \mathbf{t}_0)$, where $\mathbf{t}_0 = (\mathbf{0}, \dots, \mathbf{0})$, and let $p_1 = \mu^{(d)}(\mathbf{0}, \mathbf{t}_1)$, where $\mathbf{t}_1 = (\mathbf{e}_1, \dots, \mathbf{e}_d)$ and $\mathbf{e}_j$ is the vector in $\mathbb{Z}_2^d$ that is 0 everywhere except at component j , where it is 1. We claim that

$$(\mathbf{D}_{\mathbf{t}_0}g)(\mathbf{0}) = \prod_{S \subseteq [d]} (-1)^{\prod_{i=1}^d (\mathbf{0}_i + \sum_{j \in S} \mathbf{0}_i)} = \prod_{S \subseteq [d]} (-1)^0 = 1,$$

and

$$(\mathbf{D}_{\mathbf{t}_1}g)(\mathbf{0}) = \prod_{S \subseteq [d]} (-1)^{\prod_{i=1}^d (\mathbf{0}_i + \sum_{j \in S} \mathbf{e}_{j,i})} = (-1)^1 \cdot \prod_{S \subsetneq [d]} (-1)^0 = -1.$$

The first is clear. To see the second note that, whenever S is a proper subset of $[d]$, the factor $\mathbf{0}_i + \sum_{j \in S} \mathbf{e}_{j,i}$ in the exponent vanishes at each $i \in [d] \setminus S$, and whenever $S = [d]$, the factor $\mathbf{0}_i + \sum_{j \in S} \mathbf{e}_{j,i}$ is 1 at each $i \in [d]$.

From this,

$$\begin{aligned} \|g\|_{U^d, \mu}^{2d} &= \mathbb{E}_{(x, \mathbf{t}) \sim \mu^{(d)}} [(\mathbf{D}_{\mathbf{t}}g)(x)] = \left| \sum_{(x, \mathbf{t})} \mu^{(d)}(x, \mathbf{t}) (\mathbf{D}_{\mathbf{t}}g)(x) \right| \\ &\leq |p_0 \cdot (\mathbf{D}_{\mathbf{t}_0}g)(\mathbf{0}) + p_1 \cdot (\mathbf{D}_{\mathbf{t}_1}g)(\mathbf{0})| + \sum_{\substack{(x, \mathbf{t}): \\ (x, \mathbf{t}) \neq (\mathbf{0}, \mathbf{t}_0) \\ (x, \mathbf{t}) \neq (\mathbf{0}, \mathbf{t}_1)}} \mu^{(d)}(x, \mathbf{t}) \cdot |(\mathbf{D}_{\mathbf{t}}g)(x)| \\ &\leq |p_0 - p_1| + 1 - p_0 - p_1, \end{aligned}$$

where the first inequality follows from the triangle inequality, and the second inequality follows from the computations above and the fact that each $(D_{\mathbf{t}}g)(x)$ has unit magnitude.

Next we argue that if μ is the uniform distribution over $\mathbb{Z}_2^d$, then $\mu^{(i)}$ is the uniform distribution over $(\mathbb{Z}_2^d)^{i+1}$. We show, by induction on i , that $\mu^{(i)}(\mathbf{x}, \mathbf{t}_1, \dots, \mathbf{t}_i) = 2^{-d(i+1)}$ for every $(\mathbf{x}, \mathbf{t}_1, \dots, \mathbf{t}_i) \in (\mathbb{Z}_2^d)^{i+1}$. Since $\mu^{(0)} = \mu$, the claim is clear for $i = 0$. For $i > 0$, we have

$$\mu^{(i)}(\mathbf{x}, \mathbf{t}_1, \dots, \mathbf{t}_i) = \frac{\mu^{(i-1)}(\mathbf{x}, \mathbf{t}_1, \dots, \mathbf{t}_{i-1}) \cdot \mu^{(i-1)}(\mathbf{x} + \mathbf{t}_i, \mathbf{t}_1, \dots, \mathbf{t}_{i-1})}{\sum_{\mathbf{z}} \mu^{(i-1)}(\mathbf{z}, \mathbf{t}_1, \dots, \mathbf{t}_{i-1})}.$$

Applying the induction hypothesis, this is

$$\frac{2^{-di} \cdot 2^{-di}}{2^d \cdot 2^{-di}} = 2^{-d(i+1)}.$$

Thus, $p_0 = p_1 = 2^{-d(d+1)}$, and we get $|p_0 - p_1| + 1 - p_0 - p_1 = 1 - 2^{-d(d+1)+1}$. Putting it all together, we conclude that

$$\|g\|_{U^d, \mu} \leq (1 - 2^{-d(d+1)+1})^{1/2^d} \leq e^{-2^{-d(d+1)+1} \cdot 2^{-d}} = e^{-2^{-d^2-2d+1}}. \quad \square$$

With the ϵ -bound from Lemma 3.10 in hand, we can mimic the proof of Theorem 8.2 in Kolaitis and Kopparty [2009b] to get the proof of Lemma 3.9. First we note that the bound in the conclusion of Theorem 4.8 in Kolaitis and Kopparty [2009b] for the special case in which μ is the uniform distribution is really $(1 - \epsilon)^r$ for the ϵ from Lemma 3.10. This translates into the same bound for the conclusions of Lemmas 4.1 and 3.3 in Kolaitis and Kopparty [2009b]. The bound in the conclusion of Lemma 3.3 in Kolaitis and Kopparty [2009b] is eventually plugged into the hypothesis of Vazirani's XOR Lemma, which returns the bound $2^\ell \cdot (1 - \epsilon)^r$ on the statistical distance, where in our case, ℓ is the number of $\leq k$ -labeled label-connected label-dependent graphs with at most c unlabeled vertices. A generous bound on this factor is $\ell \leq 2^{(c+k)^2+1}$. Now note that in the proof of Theorem 8.2 in Kolaitis and Kopparty [2009b], the remaining parameters are $r = \Omega_k(n/c)$ and $d = O(c^2)$. Using the bound on ϵ from Lemma 3.10, we get the following bound on the statistical distance:

$$2^\ell \cdot (1 - \epsilon)^r \leq 2^{(c+k)^2+1} \cdot (\exp(-2^{-O(c^4)}))^{\Omega_k(n/c)}.$$

For $c \leq \log \log \log n$, this is $2^{-\Omega_k(n/\log n)}$, as required.

3.5. The Argument Itself

Finally, we are at the point where we can execute the plan sketched at the beginning of Section 3. Fix a positive integer k (for the application in Section 5, it suffices to take $k = 3$) and let $p(x_1, \dots, x_k)$ be a regular normal form of order $c \leq \log \log \log n$. For every $\mathbf{a} = (a_1, \dots, a_k) \in [n]^k$, define the following indicator random variables:

$$\begin{aligned} X(\mathbf{a}) &:= \mathbb{I}[p(\mathbf{a}) \not\leftrightarrow p(\mathbf{a} \circ \pi) \text{ for some } \pi \in S_k], \\ Y(\mathbf{a}) &:= \mathbb{I}[p(\mathbf{a}) \leftrightarrow p(\mathbf{a} \circ \pi) \text{ for every } \pi \in S_k]. \end{aligned}$$

Obviously, $X(\mathbf{a}) = 1 - Y(\mathbf{a})$, and $Y(\mathbf{a})$ is the indicator random variable for the event that p does not distinguish any two permuted versions of $\mathbf{a}$. Our goal is to show that in a random graph G , with high probability, $Y(\mathbf{a})$ holds for some $\mathbf{a}$, and for this, we will follow the plan sketched at the beginning of this section.

Write $p(\mathbf{x})$ as a DNF on the (Boolean) variables

$$\tau_1(\mathbf{x}), \dots, \tau_r(\mathbf{x}), \oplus H_1(\mathbf{x}), \dots, \oplus H_\ell(\mathbf{x}),$$

where $\tau_1, \dots, \tau_r$ are the atomic types on $x_1, \dots, x_k$, and $H_1, \dots, H_\ell$ are the $\leq k$ -labeled label-connected, label-dependent graphs with labeled vertices within $x_1, \dots, x_k$ and order at most c . Observe that, in particular, $H_1, \dots, H_\ell$ have at most c unlabeled vertices and thus belong to $\text{Conn}_{\leq k}^{*,c}$. Let us assume that $H_1, \dots, H_e$ are the ones for which $\text{aut}(H_i)$ is odd, and that $H_{e+1}, \dots, H_\ell$ are the rest. Also assume that $H_1, \dots, H_f$ are the graphs from among $H_1, \dots, H_e$ that do not have any label at all, and $H_{f+1}, \dots, H_e$ are the rest. Since atomic types are mutually exclusive and each $\oplus H_j(\mathbf{a})$ is false for $j \in \{e+1, \dots, \ell\}$, we may assume that each term in the DNF formula has the form

$$\tau_i(\mathbf{x}) \cdot \prod_{j \in K} \oplus H_j \cdot \prod_{j \in K'} \overline{\oplus H_j} \cdot \prod_{j \in I} \oplus H_j(\mathbf{x}) \cdot \prod_{j \in I'} \overline{\oplus H_j(\mathbf{x})}. \quad (8)$$

for some $i \in [r]$, some partition (K, K') of $[f]$, and some partition (I, I') of $[e] \setminus [f]$.

Next note that for every permutation $\pi \in S_k$, the sequence of Boolean variables $\tau_1(\mathbf{x} \circ \pi), \dots, \tau_r(\mathbf{x} \circ \pi)$ is equivalent to a permutation of the sequence of Boolean variables $\tau_1(\mathbf{x}), \dots, \tau_r(\mathbf{x})$. Similarly, the sequence of Boolean variables $\oplus H_{f+1}(\mathbf{x} \circ \pi), \dots, \oplus H_e(\mathbf{x} \circ \pi)$ is equivalent to a permutation of the sequence $\oplus H_{f+1}(\mathbf{x}), \dots, \oplus H_e(\mathbf{x})$. Therefore, $p(\mathbf{x})$ and $p(\mathbf{x} \circ \pi)$ are functions of the same Boolean variables, and we can write $p(\mathbf{x} \circ \pi)$ also as a DNF formula with terms of type Eq. (8).

From now on, for every $K \subseteq [f]$, let R_K be the term

$$R_K := \prod_{j \in K} \oplus H_j \cdot \prod_{j \in K'} \overline{\oplus H_j},$$

where $K' = [f] \setminus K$. Recall that $H_1, \dots, H_f$ are all label-free, and therefore R_K does not depend on $\mathbf{x}$. Similarly, for every $I \subseteq [e] \setminus [f]$, let $S_I(\mathbf{x})$ be the term

$$S_I(\mathbf{x}) := \prod_{j \in I} \oplus H_j(\mathbf{x}) \cdot \prod_{j \in I'} \overline{\oplus H_j(\mathbf{x})},$$

where $I' = ([e] \setminus [f]) \setminus I$. For every $K \subseteq [f]$, let $p_K(\mathbf{x})$ denote the disjunction of the terms in $p(\mathbf{x})$ that are consistent with R_K . Therefore, $p(\mathbf{x})$ is equivalent to the disjunction $\bigvee_{K \subseteq [f]} p_K(\mathbf{x})$.

Define the *all-positive-term* as one in which $\oplus H_j(\mathbf{x})$ appears positively for all $j \in [e] \setminus [f]$:

$$Z_K(\mathbf{x}) := \sigma(\mathbf{x}) \cdot R_K \cdot S_{[e] \setminus [f]}(\mathbf{x}),$$

where σ is the atomic type that forces $x_i \neq x_j$ for $i \neq j$, and all possible edges among different x_i, x_j . We show that for every $\mathbf{a} \in [n]^k$, the event $Z_K(\mathbf{a}) = 1$ implies $p(\mathbf{a}) \leftrightarrow p(\mathbf{a} \circ \pi)$ for every $\pi \in S_k$.

LEMMA 3.11. $Z_K(\mathbf{a}) \leq Y(\mathbf{a})$.

PROOF. Fix a permutation $\pi \in S_k$. First note that the choice of σ guarantees that $\sigma(\mathbf{a})$ is equivalent to $\sigma(\mathbf{a} \circ \pi)$. Also, the sequence $\oplus H_{f+1}(\mathbf{a} \circ \pi), \dots, \oplus H_e(\mathbf{a} \circ \pi)$ is equivalent to a permutation of the sequence $\oplus H_{f+1}(\mathbf{a}), \dots, \oplus H_e(\mathbf{a})$, and all appear positively in $Z_K(\mathbf{a})$. It follows from these facts that either the term $Z_K(\mathbf{a})$ appears in both DNFs for $p(\mathbf{a})$ and $p(\mathbf{a} \circ \pi)$, or in neither. If it appears in both, then clearly $Z_K(\mathbf{a}) = 1$ implies both $p(\mathbf{a})$ and $p(\mathbf{a} \circ \pi)$. If it does not appear in either, then $Z_K(\mathbf{a}) = 1$ implies $p(\mathbf{a})$ and $p(\mathbf{a} \circ \pi)$, since $Z_K(\mathbf{a})$ is incompatible with any other term of the DNFs for $p(\mathbf{a})$ and $p(\mathbf{a} \circ \pi)$. In either case, $Z_K(\mathbf{a})$ implies $p(\mathbf{a}) \leftrightarrow p(\mathbf{a} \circ \pi)$. $\square$

At this point, it will suffice to show that for every $K \subseteq [f]$, the event $Z_K(\mathbf{a}) = 1$ holds for some $\mathbf{a} \in [n]^k$ with high probability in the probability space conditioned on R_K . From now on, for every event A , write

$$\mathbb{P}_K[A] := \mathbb{P}[A \mid R_K].$$

Let us start by computing the probability of $Z_K(\mathbf{a})$ for $\mathbf{a} \in [n]^k$ with $a_i \neq a_j$ for $i \neq j$ in this probability space. Let δ be the maximum, over all injective atomic types $\tau(\mathbf{x})$, of the statistical distance between the distribution $\text{freq}_{\leq k, G}^{*,c}(\mathbf{a})$ as $G \sim G(n, 1/2 \mid \tau(\mathbf{a}))$ and the uniform distribution over $\text{FFreq}^*(\tau, \leq k, c)$. Note that, by symmetry, δ does not depend on $\mathbf{a}$ provided $a_i \neq a_j$ for $i \neq j$.

LEMMA 3.12.

$$\mathbb{P}_K[Z_K(\mathbf{a})] \leq \left(\frac{2^{-e}}{2^{-f}} + \delta \cdot \frac{1}{2^{-f} - \delta} + \delta \cdot \frac{2^{-e}}{2^{-f} \cdot (2^{-f} - \delta)} \right) \cdot 2^{-\binom{k}{2}};$$

and

$$\mathbb{P}_K[Z_K(\mathbf{a})] \geq \left(\frac{2^{-e}}{2^{-f}} - \delta \cdot \frac{1}{2^{-f} + \delta} - \delta \cdot \frac{2^{-e}}{2^{-f} \cdot (2^{-f} + \delta)} \right) \cdot 2^{-\binom{k}{2}}.$$

PROOF. We have

$$\mathbb{P}_K[Z_K(\mathbf{a})] = \frac{\mathbb{P}[Z_K(\mathbf{a}) \cdot R_K]}{\mathbb{P}[R_K]} = \frac{\mathbb{P}[S_{[e] \setminus [f]}(\mathbf{a}) \cdot R_K \mid \sigma(\mathbf{a})] \cdot \mathbb{P}[\sigma(\mathbf{a})]}{\mathbb{P}[R_K]}.$$

The denominator is at most $2^{-f} + \delta$ and at least $2^{-f} - \delta$ by choice of δ . Similarly, the numerator is at most $(2^{-e} + \delta) \cdot 2^{-\binom{k}{2}}$ and at least $(2^{-e} - \delta) \cdot 2^{-\binom{k}{2}}$ also by choice of δ . Note that $\mathbb{P}[\sigma(\mathbf{a})] = 2^{-\binom{k}{2}}$. Now,

$$\frac{2^{-e} + \delta}{2^{-f} - \delta} - \frac{2^{-e}}{2^{-f}} = \frac{2^{-e} \cdot 2^{-f} + \delta \cdot 2^{-f} - 2^{-e} \cdot 2^{-f} + 2^{-e} \cdot \delta}{2^{-f} \cdot (2^{-f} - \delta)}$$

which simplifies to

$$\delta \cdot \frac{1}{2^{-f} - \delta} + \delta \cdot \frac{2^{-e}}{2^{-f} \cdot (2^{-f} - \delta)}.$$

Similarly,

$$\frac{2^{-e}}{2^{-f}} - \frac{2^{-e} - \delta}{2^{-f} + \delta} = \delta \cdot \frac{1}{2^{-f} + \delta} + \delta \cdot \frac{2^{-e}}{2^{-f} \cdot (2^{-f} + \delta)}.$$

□

Next we compute, for every $\mathbf{a}, \mathbf{a}' \in [n]^k$ with all $a_1, \dots, a_k, a'_1, \dots, a'_k$ distinct, the probability of $Z_K(\mathbf{a}) \cdot Z_K(\mathbf{a}')$ in the probability space conditioned on R_K . Let γ be the maximum, over all injective atomic types $\tau(\mathbf{x}, \mathbf{x}')$, of the statistical distance between the distribution $\text{freq}_{\leq 2k, G}^{*,c}(\mathbf{a}, \mathbf{a}')$ as $G \sim G(n, 1/2 \mid \tau(\mathbf{a}, \mathbf{a}'))$ and the uniform distribution over $\text{FFreq}^*(\tau, \leq 2k, c)$. Note that, by symmetry, γ does not depend on $\mathbf{a}, \mathbf{a}'$ provided they are all distinct.

LEMMA 3.13.

$$\mathbb{P}_K[Z_K(\mathbf{a}) \cdot Z_K(\mathbf{a}')] \leq \left(\frac{2^{-2e}}{2^{-2f}} + \gamma \cdot \frac{1}{2^{-f} - \gamma} + \gamma \cdot \frac{2^{-2e+f}}{2^{-f} \cdot (2^{-f} - \gamma)} \right) \cdot 2^{-2\binom{k}{2}}.$$

PROOF. Let A denote the event that $\sigma(\mathbf{a})$ and $\sigma(\mathbf{a}')$ both hold. We have

$$\begin{aligned}\mathbb{P}_K[Z_K(\mathbf{a}) \cdot Z_K(\mathbf{a}')] &= \frac{\mathbb{P}[Z_K(\mathbf{a}) \cdot Z_K(\mathbf{a}') \cdot R_K]}{\mathbb{P}[R_K]} \\ &= \frac{\mathbb{P}[S_{[e] \setminus [f]}(\mathbf{a}) \cdot S_{[e] \setminus [f]}(\mathbf{a}') \cdot R_K \mid A] \cdot \mathbb{P}[A]}{\mathbb{P}[R_K]}.\end{aligned}$$

The denominator is at least $2^{-f} - \gamma$ by choice of γ . The numerator is at most $(2^{-2e+f} + \gamma) \cdot 2^{-2\binom{k}{2}}$ also by choice of γ . The trailing $2^{-2\binom{k}{2}}$ factor is $\mathbb{P}[A]$. Now:

$$\frac{2^{-2e+f} + \gamma}{2^{-f} - \gamma} - \frac{2^{-2e+f}}{2^{-f}} = \frac{2^{-2e+f} \cdot 2^{-f} + \gamma \cdot 2^{-f} - 2^{-2e+f} \cdot 2^{-f} + 2^{-2e+f} \cdot \gamma}{2^{-f} \cdot (2^{-f} - \gamma)}$$

which simplifies to

$$\gamma \cdot \frac{1}{2^{-f} - \gamma} + \gamma \cdot \frac{2^{-2e+f}}{2^{-f} \cdot (2^{-f} - \gamma)}.$$

Let us note at this point that the number of $\leq k$ -labeled graphs of order at most c is bounded by 2^{c^2+1} . Therefore, using the bound $c \leq \log \log \log n$, we have $\ell \leq \frac{1}{2} \log n$ for sufficiently large n , and in particular

$$2^\ell \leq \sqrt{n}. \quad (9)$$

We use this to prove the main consequence of this analysis up to now.

LEMMA 3.14. *Let $\mathbf{a}, \mathbf{a}' \in [n]^k$ be such that $a_1, \dots, a_k, a'_1, \dots, a'_k$ are all different. The following hold.*

- (1) $\mathbb{P}_K[Z_K(\mathbf{a})] \geq n^{-1/2} \cdot 2^{-\binom{k}{2}} - 2^{-\Omega_k(n/\log n)}$.
- (2) $|\mathbb{P}_K[Z_K(\mathbf{a}) \cdot Z_K(\mathbf{a}')] - \mathbb{P}_K[Z_K(\mathbf{a})] \cdot \mathbb{P}_K[Z_K(\mathbf{a}')]| \leq 2^{-\Omega_k(n/\log n)}$.

PROOF. By Lemma 3.9, both δ and γ are $2^{-\Omega_k(n/\log n)}$. On the other hand, we have $2^{e-f} \leq 2^e \leq 2^\ell \leq n^{1/2}$ by Eq. (9) and also $2^f \leq 2^\ell \leq n^{1/2}$ by (9). Therefore, $2^{f-e} \geq 2^{-e} \geq 2^{-\ell} \geq n^{-1/2}$ and $2^{-f} \geq 2^{-\ell} \geq n^{-1/2}$. Now (1) follows from plugging these bounds into the lower bound in Lemma 3.12 and (2) follows from plugging these bounds into the upper bound in Lemma 3.12 and the bound in Lemma 3.13 and recalling that k is a constant. $\square$

Now we conclude by proving the main result of this section.

LEMMA 3.15. *For every $k > 0$ and $\epsilon > 0$, there exists $n_0 \geq 0$ such that for every $n \geq n_0$ and every regular form $p(x_1, \dots, x_k)$ of order bounded by $\log \log \log n$, for $G \sim G(n, 1/2)$, the probability that there exists $\mathbf{a} \in [n]^k$ with $a_i \neq a_j$ for $i \neq j$ such that $p(\mathbf{a}) \leftrightarrow p(\mathbf{a} \circ \pi)$ for every $\pi \in S_k$ is at least $1 - \epsilon$.*

PROOF. Fix k and ϵ and choose n_0 large. Let $m = \lfloor n/k \rfloor$. Divide $[n]$ into m disjoint k -tuples $(\mathbf{a}_1, \dots, \mathbf{a}_m)$ arbitrarily but in such a way that $a_{\ell,i} \neq a_{\ell,j}$ for $i \neq j$. Define $Y = \sum_{\ell \in [m]} Y(\mathbf{a}_\ell)$ and $Z_K = \sum_{\ell \in [m]} Z_K(\mathbf{a}_\ell)$, the second for every $K \subseteq [f]$. Note that by Lemma 3.11, we have $Z_K \leq Y$. We want to show that $\mathbb{P}_K[Z_K = 0] \leq \epsilon$. This will be enough, since then,

$$\mathbb{P}[Y = 0] = \sum_{K \subseteq [e]} \mathbb{P}[Y = 0 \mid R_K] \cdot \mathbb{P}[R_K] \leq \sum_{K \subseteq [e]} \mathbb{P}_K[Z_K = 0] \cdot \mathbb{P}[R_K] \leq \epsilon.$$

To show that $\mathbb{P}_K[Z_K = 0] \leq \epsilon$, we proceed by the second moment method. To simplify notation, let us fix $K \subseteq [f]$ and abbreviate Z_K by Z , and $Z_K(\mathbf{a}_\ell)$ by Z_ℓ . Similarly, all

expectations $\mathbb{E}$, variances $\mathbb{V}$, and probabilities $\mathbb{P}$ appearing next refer to the probability space $\mathbb{P}_K$. In computing the variance $\mathbb{V}[Z] = \mathbb{E}[Z^2] - \mathbb{E}[Z]^2$, we have

$$\begin{aligned}
 \mathbb{E}[Z^2] - \mathbb{E}[Z]^2 &= \sum_{i,j} \mathbb{E}[Z_i \cdot Z_j] - \sum_{i,j} \mathbb{E}[Z_i] \cdot \mathbb{E}[Z_j] \\
 &\leq \sum_i \mathbb{E}[Z_i^2] + 2 \cdot \sum_{i \neq j} (\mathbb{E}[Z_i \cdot Z_j] - \mathbb{E}[Z_i] \cdot \mathbb{E}[Z_j]) \\
 &\leq \sum_i \mathbb{E}[Z_i] + 2 \cdot \sum_{i \neq j} (\mathbb{P}[Z_i \cdot Z_j] - \mathbb{P}[Z_i] \cdot \mathbb{P}[Z_j]) \\
 &= \mathbb{E}[Z] + 2 \cdot \binom{m}{2} \cdot 2^{-\Omega_k(n/\log n)} \\
 &= \mathbb{E}[Z] + 2^{-\Omega_k(n/\log n)},
 \end{aligned}$$

where the first inequality follows from considering the case $i = j$ in the first double sum and ignoring it in the second, the next inequality follows from the fact that Z_i is a 0-1-random variable, the equality after it follows from Lemma 3.14.2 (recall that $\mathbb{P}$ really stands for $\mathbb{P}_K$ here), and the last equality follows from $m = \lfloor n/k \rfloor$ and the fact that k is a constant.

Now by Lemma 3.14.1, we have

$$\mathbb{E}[Z] \geq m \cdot (n^{-1/2} \cdot 2^{-\binom{k}{2}} - 2^{-\Omega_k(n/\log n)}) = \Omega_k(n^{1/2}).$$

Applying it to Chebyshev's inequality, we obtain

$$\mathbb{P}[Z = 0] \leq \frac{\mathbb{V}[Z]}{\mathbb{E}[Z]^2} \leq \frac{\mathbb{E}[Z] + 2^{-\Omega_k(n/\log n)}}{\mathbb{E}[Z]^2} \leq \epsilon$$

for sufficiently large n . $\square$

4. DEFINING A LINEAR PRE-ORDER OF WIDTH TWO

In this section, we construct the formula of very low depth that defines a linear pre-order of width 2 with high probability. The proof strategy is to analyze a variant of an algorithm for graph canonization due to Karp [1979], and to exploit its massive implicit parallelism to get formulas of very low depth.

4.1. Plan of Action

Informally, the graph canonization algorithm works as follows. For a given graph G , split the vertices into two classes: those of even degree and those of odd degree. Inductively, we split the classes further by dividing the vertices according to the parity of the numbers of neighbours they have in each of the existing classes. We continue this process until no more classes are split.

We will need three facts about this process: (1) that for $G \sim G(n, 1/2)$ the process will reach a state where each class has at most two vertices with high probability, (2) that this will happen in fewer than n “generations” of the splitting process with high probability, and (3) that the process is massively parallel: all the classes created between the $\ell/2$ -th generation and the ℓ -th generation are definable in terms of the classes created in the $(\log_2 \ell)$ -th generation.

4.2. Splitting Procedure

Let $G = (V, E)$ be an undirected graph. For a vertex x and a set B , we write $p(G, x, B)$ for the parity of the number of neighbours that x has in B . We extend this to the

following set.

$$p(G, A, B) = \sum_{x \in A} p(G, x, B) \pmod{2}.$$

A *splitting tree* for G is a rooted binary tree T with each node t carrying a label $L_t \subseteq V$ and a sign $M_t \in \{+, -\}$ denoting whether t is *marked* or *unmarked*, and satisfying the following properties:

- (1) the label of the root is V ,
- (2) no two siblings are marked,
- (3) if t is an internal node, then¹ $L_{t0} \cup L_{t1} = L_t$ and $L_{t0} \cap L_{t1} = \emptyset$,
- (4) if s is a leaf, $x, y \in L_s$ and t is marked, then $p(G, x, L_t) = p(G, y, L_t)$.

Given a splitting tree T for G , let $R(T)$ denote the set of unmarked nodes that are either the root or are a left child. Let $R'(T)$ be the subset of $R(T)$ containing the root and all nodes t such that the label of t and its sibling are both nonempty.² One step of the splitting procedure works as follows.

- (1) Let t be the least node in $R(T)$ in level-order³ and mark it.
- (2) For every leaf s , let $L_{sa} := \{x \in L_s : p(G, x, L_t) = a\}$ for both $a = 0$ and $a = 1$.
- (3) Make⁴ s_0 and s_1 the left and right children of s and leave them unmarked.

Let $\mathcal{P}(T)$ be the result of applying one step of the splitting procedure to T . If the node t that is chosen in the first step also belongs to $R'(T)$, we say that the step is *proper*, otherwise *improper*. When $R'(T)$ is empty, we say that the procedure *stalls* at T . Note that when it stalls, it will never make a proper step again. The procedure starts at the splitting tree T_0 that has only an unmarked root labeled by V .

4.3. Analysis of the Splitting Procedure

Let T_0 be the tree that has only an unmarked root labeled by V . For $k \geq 1$, let $T_k := \mathcal{P}(T_{k-1})$. Ideally, we would like to show that after a modest number of steps, all leaves of the splitting tree are labeled by singletons or empty sets. Unfortunately, the splitting procedure is not able to produce a tree with this property in general, not even with high probability on a random graph. The best we will be able to show is that for a randomly generated graph, with high probability, all leaves will have at most two vertices.

We identify three key desirable properties of T_k , where the third is our goal:

- (A_k). T_k has $L_t \neq \emptyset$ for every node t ,
- (B_k). T_k has been generated through proper steps only,
- (C_k). T_k has $|L_t| \leq 2$ for every leaf t .

Next, we will show the following.

- (1) Property (A_k) holds with high probability for suitable values of k .
- (2) Property (A_k) implies (B_{2k}) for every graph and every $k \geq 0$.
- (3) Conditioned on (B_{2k}), property (C_{2k}) holds with high probability for suitable values of k .

¹Karp requires also $L_{t0} \neq \emptyset$ and $L_{t1} \neq \emptyset$. For us, it is convenient to not require it, and Karp's analysis still goes through with minor modifications that we point out.

²Karp defines $R(T)$ as the set of unmarked nodes t that are either the root or that have a sibling t' such that $|L_{t'}| > |L_t|$, or $|L_{t'}| = |L_t|$ and are a left child. This difference is inessential to the analysis. The only important point is to unambiguously choose one of the two children when both are unmarked and nonempty.

³Karp used symmetric order. This difference is not essential for Karp's analysis but is important for us.

⁴Karp's version makes this step only if $L_{sa} \neq \emptyset$ for both $a = 0$ and $a = 1$; this note is related to footnote 1.

Before we analyse the probability of (A_k) , we need to introduce some terminology and a lemma from Karp [1979]. Let T be a splitting tree for some graph H on the vertices V . To every node $t \in R'(T)$, we associate a set $S_t \subseteq V$: the set of all x for which t is the maximal node in $R'(T)$ such that L_t contains x . Let $\mathcal{S}(T)$ be the collection of all such sets. For every $t \in R'(T)$, let β_t be t together with the set of nodes $s \in R'(T)$ such that $s \neq t$ and L_s is a maximal subset of L_t . Note that $S_t = \Delta_{s \in \beta_t} L_s$, where Δ denotes symmetric difference. Define $\ell(x, S_t) := \sum_{s \in \beta_t} p(H, x, L_s) \bmod 2$. We will say that another graph G on the vertices V is consistent with T if $p(G, x, L_t) = p(H, x, L_t)$ holds for every $x \in V$ and every node $t \in R'(T)$.

We state a consequence of Lemmas 4 and 5 in Karp [1979].⁵

LEMMA 4.1. *Let T be the splitting tree of some graph on the vertices V and let H be chosen uniformly at random among the graphs on the vertices V that are consistent with T . If t is a node in $R'(T)$, then the distribution of $\{p(H, x, L_t)\}_{x \in V}$ is uniform over the assignments that satisfy the constraints*

$$p(H, S, L_t) = \ell(L_t, S) \text{ for every } S \in \mathcal{S}(T_k) \setminus \{Y\},$$

where Y is the unique set in $\mathcal{S}(T)$ of which L_t is a proper subset.

In order to be able to make use of this lemma, it is important to notice that if G denotes a random graph drawn from $G(n, 1/2)$ and $T_0, T_1, \dots$ denotes the random sequence of splitting trees produced by this random graph, then the distribution of T_{k+1} conditioned on T_k is equally produced as follows: first choose a graph H uniformly at random among those consistent with T_k , and then run one step of the splitting procedure on T_k with respect to H . This follows from the fact that the restriction of a uniform distribution to a subset of its support is uniformly distributed on that subset.

Now we can analyze the probability of (A_k) .

LEMMA 4.2. *Let $n \geq 1$ and $k \geq 1$ be integers such that $4k \leq \log_2 n$, and let $G \sim G(n, 1/2)$. Then, the probability that (A_k) fails is $2^{k+1} \cdot \exp(-n/2^{6k})$.*

PROOF. In order to simplify notation, in this proof, we let $n_t := |L_t|$. For a node t at depth $\ell \leq k$ in T_k , we say that t is *unbiased* if $|n_t - n \cdot 2^{-\ell}| \leq n \cdot 2^{-(2k-\ell+1)}$ holds, and *biased* otherwise. Note for later use that we allow the error-term $n \cdot 2^{-(2k-\ell+1)}$ to grow with ℓ , but that it always stays below $n \cdot 2^{-\ell}$ because $\ell \leq k$. Let us consider the event defined as follows.

(A'_k) . T_k has every node unbiased.

Note that since the error-term for $\ell = k$ is smaller than $n \cdot 2^{-k}$, property (A'_k) implies (A_k) . Thus, it suffices to bound the probability that (A'_k) fails.

Since (A'_0) holds, if (A'_k) fails, then there is a largest $\ell \in \{0, \dots, k-1\}$ such that (A'_ℓ) is true and $(A'_{\ell+1})$ is false. Fix $\ell \in \{0, \dots, k-1\}$, a leaf r of T_ℓ , and $a \in \{0, 1\}$, and we bound the probability that the child ra of r becomes biased in $T_{\ell+1}$ conditioned on T_ℓ satisfying (A'_ℓ) . Let t be the node with respect to which the splitting step $\ell+1$ is made. Since we are assuming that T_ℓ satisfies (A'_ℓ) , each label is nonempty, and therefore, t belongs to $R'(T_\ell)$. Let Y be the unique set in $\mathcal{S}(T_\ell)$ of which L_t is a proper subset. By the discussion after Lemma 4.1, the tree $T_{\ell+1}$ can be seen as produced by first choosing H uniformly at random among the graphs that are consistent with T_ℓ , and then applying the splitting procedure on T_ℓ with respect to t and H . By Lemma 4.1, the distribution

⁵It would seem from Lemma 4 in Karp [1979] that we also need the constraint $p(G, L_t, L_t) = 0$. However, in our notation this constraint is implicit, since $p(G, L_t, L_t)$ counts each edge within L_t exactly twice.

of $\{p(H, x, L_t)\}_{x \in V}$ is uniform over the assignments that satisfy the constraints

$$p(H, S, L_t) = \ell(L_t, S) \quad (10)$$

for every S in $S(T_\ell) \setminus \{Y\}$. In particular, since all sets in $S(T_\ell)$ are pairwise disjoint, if S is the unique minimal set in $S(T_\ell)$ that contains L_r , then the distribution of $\{p(H, x, L_t)\}_{x \in S}$ is uniform over the assignments that satisfy Constraint (10) for this S only, or no constraint at all if $S = Y$.

Since r is unbiased, the set L_r is nonempty. Fix $x_0 \in L_r \subseteq S$. A different way of generating the distribution $\{p(H, x, L_t)\}_{x \in S}$ without sampling H is by first choosing values for $p(-, x, L_t)$ for $x \in S \setminus \{x_0\}$ uniformly and independently at random, and then setting the value for $p(-, x_0, L_t)$ to the unique value that satisfies the constraint $p(-, S, L_t) = \ell(L_t, S)$, or setting it uniformly and independently at random if $S = Y$. In either case, the number X of elements x in $L_r \setminus \{x_0\}$ for which $p(-, x, L_t) = a$ is a random variable distributed according to the binomial distribution $B(m, \frac{1}{2})$ with $m = n_r - 1$. Note for later use that $|n_{ra} - X| \leq 1$, because only x_0 could be missed in the count. By Hoeffding's inequality for the binomial distribution, the probability that $|X - \frac{1}{2} \cdot m| \geq t$ is bounded by $2e^{-2t^2/m}$, which is bounded by

$$2e^{-2t^2/(2n \cdot 2^{-\ell})}, \quad (11)$$

because r is unbiased, and hence $m = n_r - 1 < n \cdot 2^{-\ell} + n \cdot 2^{-(2k-\ell+1)} \leq 2n \cdot 2^{-\ell}$ because $\ell < k$. Now, if ra were biased, we would have

$$|n_{ra} - n \cdot 2^{-(\ell+1)}| > n \cdot 2^{-(2k-(\ell+1)+1)}.$$

Since $|n_{ra} - X| \leq 1$ and since $|\frac{1}{2} \cdot n_r - n \cdot 2^{-(\ell+1)}| \leq \frac{1}{2} \cdot n \cdot 2^{-(2k-\ell+1)}$ because r is unbiased, by the triangle inequality, this would mean that

$$\left| X - \frac{1}{2} \cdot n_r \right| > n \cdot 2^{-(2k-(\ell+1)+1)} - \frac{1}{2} \cdot n \cdot 2^{-(2k-\ell+1)} - 1,$$

and in particular, using $0 \leq \ell \leq k-1$ and $4k \leq \log_2 n$, that

$$\left| X - \frac{1}{2} \cdot (n_r - 1) \right| \geq n \cdot 2^{-3k}.$$

The probability of this happening is bounded by Eq. (11) with $t = n \cdot 2^{-3k}$, which is at most $\epsilon := 2 \exp(-n \cdot 2^{-6k})$.

The argument is now finished by two union bounds. By the union bound over the 2^ℓ leaves of T_ℓ , the probability that some leaf of T_ℓ generates a biased child is at most $2^\ell \cdot \epsilon$. By the union bound over ℓ , the probability that there exists an $\ell \in \{0, \dots, k-1\}$ for which (A'_ℓ) holds but $(A'_{\ell+1})$ fails is at most $\sum_{\ell=0}^{k-1} 2^\ell \cdot \epsilon$. Thus, the probability that (A'_k) fails is bounded by $2^{k+1} \cdot \exp(-n \cdot 2^{-6k})$. $\square$

Next we observe that (A_k) implies (B_{2^k}) .

LEMMA 4.3. *For any graph G and $k \geq 0$, if (A_k) holds, then (B_{2^k}) holds.*

PROOF. In a complete binary tree of depth k , the number of *left* children at depth at most k is $\sum_{i=1}^k 2^{i-1} = 2^k - 1$. Now, if T_k satisfies (A_k) , then the root and every left child at depth at most k has generated a proper step in the process of producing T_{2^k} ; these are 2^k proper steps as claimed. $\square$

Finally we note that 3-element sets split with high probability if enough steps are proper. This is similar to Lemma 7 in Karp [1979].

LEMMA 4.4. *Let $G \sim G(n, 1/2)$ and let $k \geq 0$. Then, the probability that (B_k) holds and (C_k) fails is at most $\binom{n}{3} \cdot 2^{-2k}$.*

PROOF. Fix a 3-element set $A \subseteq V$ and fix $\ell \leq k$. Let S_ℓ denote the event that the set A is not split at step ℓ and P_ℓ denote the event that step ℓ is proper. We aim to show that $\mathbb{P}[\bigcap_{\ell=1}^k S_\ell \cap \bigcap_{\ell=1}^k P_\ell] \leq 2^{-2k}$ and the result then follows by a union bound over all three element subsets.

Now,

$$\mathbb{P}\left[\bigcap_{\ell=1}^k S_\ell \cap \bigcap_{\ell=1}^k P_\ell\right] = \prod_{\ell=0}^{k-1} \mathbb{P}\left[S_{\ell+1} \cap P_{\ell+1} \mid \bigcap_{i=1}^{\ell} (S_i \cap P_i)\right],$$

which is bounded by

$$\prod_{\ell=0}^{k-1} \mathbb{P}\left[S_{\ell+1} \mid P_{\ell+1} \cap \bigcap_{i=1}^{\ell} (S_i \cap P_i)\right]. \quad (12)$$

So, it suffices to show that each term in Eq. (12) is bounded by $\frac{1}{4}$.

Fix $\ell \in \{0, \dots, k-1\}$ and let $\mathbf{T}$ denote the sequence of splitting trees $T_0, \dots, T_\ell$. Let $\mathcal{T}$ denote the set of all sequences of splitting trees of length $\ell+1$ and $\mathcal{T}_A$ denote the subset of $\mathcal{T}$ consisting of those sequences $\mathbf{U} = U_0, \dots, U_\ell$ in which all steps are proper and A does not split at any stage and U_ℓ splits properly, that is, with respect to a node in $R'(U_\ell)$. In other words, the sequence $\mathbf{U}$ satisfies $P_{\ell+1} \cap \bigcap_{i=1}^{\ell} (S_i \cap P_i)$. We now argue that, for any given $\mathbf{U} \in \mathcal{T}$, we have $\mathbb{P}[S_{\ell+1} \mid \mathbf{T} = \mathbf{U}] \leq \frac{1}{4}$.

Let r be a leaf of T_ℓ such that $A \subseteq L_r$. Let t be the node of T_ℓ with respect to which the splitting step $\ell+1$ is made. We argue that, conditioned on the event that this step is proper, that is, t belongs to $R'(T_\ell)$, the probability that the elements of A are not split apart in $T_{\ell+1}$ is at most $1/4$. Let Y be the unique set in $S(T_\ell)$ of which L_t is a proper subset. By the discussion after Lemma 4.1, the tree $T_{\ell+1}$ can be seen as produced by first choosing H uniformly at random among the graphs that are consistent with T_ℓ , and then applying the splitting procedure on T_ℓ with respect to t and H . By Lemma 4.1, the distribution of $\{p(H, x, L_t)\}_{x \in V}$ is uniform over the assignments that satisfy the constraints

$$p(G, S, L_t) = \ell(L_t, S), \quad (13)$$

for every $S \in S(T_\ell) \setminus \{Y\}$. In particular, since all sets in $S(T_\ell)$ are pairwise disjoint, if S is the unique minimal set in $S(T_\ell)$ that contains L_r , then the distribution of $\{p(H, x, L_t)\}_{x \in S}$ is uniform over the assignments that satisfy Constraint (13) for this S only, or no constraint at all if $S = Y$. Thus, in case $S \neq Y$, there are $2^{|S \cup L_t| - 1}$ choices for $\{p(H, x, L_t)\}_{x \in S}$ and $2^{|S \cup L_t| - 3}$ such choices that are constant over A , and in case $S = Y$, there are $2^{|S \cup L_t|}$ choices for $\{p(H, x, L_t)\}_{x \in S}$ and $2^{|S \cup L_t| - 2}$ such choices that are constant over A . In both cases, this gives probability $1/4$, as claimed.

To complete the argument, let E_ℓ denote the event $P_{\ell+1} \cap \bigcap_{i=1}^{\ell} (S_i \cap P_i)$. We have

$$\begin{aligned} \mathbb{P}[S_{\ell+1} \mid E_\ell] &= \mathbb{P}[S_{\ell+1} \cap E_\ell] \cdot \mathbb{P}[E_\ell]^{-1} \\ &= \sum_{\mathbf{U} \in \mathcal{T}} \mathbb{P}[S_{\ell+1} \cap E_\ell \mid \mathbf{T} = \mathbf{U}] \cdot \mathbb{P}[\mathbf{T} = \mathbf{U}] \cdot \mathbb{P}[E_\ell]^{-1} \\ &= \sum_{\mathbf{U} \in \mathcal{T}_A} \mathbb{P}[S_{\ell+1} \mid \mathbf{T} = \mathbf{U}] \cdot \mathbb{P}[\mathbf{T} = \mathbf{U}] \cdot \mathbb{P}[E_\ell]^{-1} \\ &\leq \frac{1}{4} \cdot \sum_{\mathbf{U} \in \mathcal{T}_A} \mathbb{P}[\mathbf{T} = \mathbf{U}] \cdot \mathbb{P}[E_\ell]^{-1} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{4} \cdot \mathbb{P}[E_\ell] \cdot \mathbb{P}[E_\ell]^{-1} \\
&= \frac{1}{4}.
\end{aligned}$$

This completes the proof of the lemma. $\square$

We are ready to synthesize what we have learned in a single lemma. In its statement, the choice of parameters is made to minimize the probability of failure. Other choices with other goals would work as well.

LEMMA 4.5. *Let $G \sim G(n, 1/2)$. Then, the probability that $T_{\lceil n^{1/5} \rceil}$ does not satisfy $(C_{\lceil n^{1/5} \rceil})$ is at most $2^{-\Omega(n^{1/6})}$.*

PROOF. Choose $k = \lceil \frac{1}{5} \log_2 n \rceil$ in Lemma 4.2 and $k = \lceil n^{1/5} \rceil$ in Lemma 4.4 and link them through Lemma 4.3. $\square$

4.4. Defining the Splitting Steps

In this section, we show that sets L_t of the splitting trees T_k are definable by formulas $\psi_t(x)$ of very low quantifier rank. First, let us recall that if the splitting step is made with respect to node t , then every leaf s splits into the following sets.

$$\begin{aligned}
L_{s0} &= \{x \in L_s : p(G, x, L_t) = 0\}, \\
L_{s1} &= \{x \in L_s : p(G, x, L_t) = 1\}.
\end{aligned}$$

Note that the nodes at depth ℓ are generated by the ℓ th splitting step. For every non-root node t in a splitting tree T , let $v_T(t)$ be the node of T that generated t . In the following, let $u(1) := 1$ and $u(\ell) := \text{bin}_2(2(\ell - 1))$ for every $\ell \geq 2$.

LEMMA 4.6. *Let G be a graph and let $k \geq \ell \geq 1$. Then, for every node t at depth ℓ in T_k , we have $v_{T_k}(t) = u(\ell)$.*

PROOF. Let us write $T = T_k$. If t is one of the two nodes at depth 1, then $v_T(t)$ is the root, which agrees with $u(1)$. Assume now that t is a node at depth $\ell \geq 2$. Let num_2 be such that $\text{num}_2(\text{bin}_2(n)) = n$ for every positive integer n . We show that $\text{num}_2(v_T(t)) = 2(\ell - 1)$. We proceed by induction on ℓ . For $\ell = 2$, we have it, since then, $v_T(t)$ is the left child of the root 10, and $\text{num}_2(10) = 2$. Now, if t is a node at depth $\ell \geq 2$ and we assume that $\text{num}_2(v_T(t)) = 2(\ell - 1)$, then for every $a \in \{0, 1\}$, we have

$$\text{num}_2(v_T(ta)) = \text{num}_2(v_T(t)) + 2 = 2(\ell - 1) + 2 = 2((\ell + 1) - 1),$$

where the first follows from the fact that the nodes at level $\ell + 1$ are generated by the next left-child following $v_T(t)$ in the level-order, and that the level-order on nodes agrees with the order of the natural numbers when they are read in binary. $\square$

Now, for $a_1, \dots, a_\ell \in \{0, 1\}$, define

$$\psi_{1a_1 \dots a_\ell}(x) := \bigwedge_{\substack{i=1 \\ a_i=1}}^{\ell} \oplus z (\psi_{u(i)}(z) \wedge E(x, z)) \wedge \bigwedge_{\substack{i=1 \\ a_i=0}}^{\ell} \neg \oplus z (\psi_{u(i)}(z) \wedge E(x, z)).$$

Note that $\psi_1(x)$ is true, since then the conjunctions are empty. We show that the $\psi_t(x)$ are the formulas we are after.

LEMMA 4.7. *Let G be a graph and let $k \geq \ell \geq 0$. Then, for every node t at depth at most ℓ in T_k , the formula $\psi_t(x)$ defines the set L_t in G .*

PROOF. For every nonleaf node t at depth $\ell - 1$, we have $v_T(ta) = u(\ell)$ for both $a = 0$ and $a = 1$ by Lemma 4.6. Therefore,

$$L_{t0} = \{x \in L_t : p(G, x, L_{u(\ell)}) = 0\},$$

$$[6pt]L_{t1} = \{x \in L_t : p(G, x, L_{u(\ell)}) = 1\}.$$

Now, if $t = 1a_1a_2 \cdots a_\ell$, then unfolding the recursion, we have that L_t is the set of vertices $x \in V$ for which $p(G, x, L_{u(i)}) = a_i$ holds for every $i \in \{1, \dots, \ell\}$. This is precisely what $\psi_t(x)$ says. $\square$

Note that the quantifier rank of $\psi_t(x)$ depends only on the depth of t . Therefore, let $q(\ell)$ be the quantifier rank of $\psi_t(x)$ for some and hence every t of depth ℓ . Note that $q(\ell)$ is monotone nondecreasing.

LEMMA 4.8. $q(\ell) = O(\log^* \ell)$.

PROOF. If t is a node at depth ℓ , the largest $u(i)$ in the definition of ψ_t is $2(\ell - 1)$. Since q is monotone nondecreasing, we have

$$q(\ell) = 1 + q(|\text{bin}_2(2(\ell - 1))|).$$

Since the length of $\text{bin}_2(2(\ell - 1))$ is $\log_2(\ell) + O(1)$, this recurrence gives $q(\ell) = O(\log^* \ell)$, as claimed. $\square$

4.5. Defining the Linear Pre-Order

Finally we are ready to prove the main lemma of this section.

LEMMA 4.9. *There exists $d > 0$ such that for every $\delta > 0$, there exists $n_0 \geq 0$ such that for every $n \geq n_0$, there is a formula $\psi(x, y)$ of quantifier rank at most $d \log^* n$ such that, for $G \sim G(n, 1/2)$, the probability that φ defines a linear pre-order of width at most 2 is at least $1 - \delta$.*

PROOF. Choose d to be the universal multiplicative constant in the $O(\log^* \ell)$ notation in Lemma 4.8. Fix $\delta > 0$ and let n_0 be large enough so that for every $n \geq n_0$, the probability in Lemma 4.5 is at most δ , and $q(n) \leq d \log^* n$. For fixed $n \geq n_0$, let $k = \lceil n^{1/\delta} \rceil$, and let $\psi(x, y)$ be the following formula.

$$\bigvee_{\substack{s, t \\ s \leq t}} \psi_s(x) \wedge \psi_t(y),$$

where s and t range over the leaves of T_k in the disjunction. If T_k has all its leaves labeled by sets of size at most two, this defines a linear pre-order of width at most two. By choice of n_0 , this happens with probability at least $1 - \delta$. Finally, by Lemma 4.8, the quantifier rank of ψ is $q(k) \leq q(n) \leq d \log^* n$. $\square$

5. ESTABLISHING THE LOWER BOUND

Here we put it all together to prove Theorem 1.1.

THEOREM 1.1. *There exists a constant $c > 0$ such that for every large enough integer q , every $\epsilon > 0$, and every large enough integer n , there exists an $\text{FO}[\oplus]$ -formula $\phi(u, v, w)$ of quantifier rank q such that, for every Boolean combination p of quantifier-free formulas and $\text{FO}[\oplus]$ -polynomials of degree bounded by a tower of exponentials of height at most q/c , the formulas ϕ and p must disagree on a fraction $1 - \epsilon$ of all graphs with n vertices.*

PROOF. Let d be the constant in Lemma 4.9 and choose $c := d + 1$. Choose q_0 large enough (to be determined later), fix $q \geq q_0$ and $\epsilon > 0$, and choose n_0 large enough (to be determined later) and fix $n \geq n_0$. Let $\psi(x, y)$ be the formula from Lemma 4.9 for $\delta = \epsilon/2$

and the fixed n ; in particular, n_0 must be larger than the n_0 from Lemma 4.9 for this particular δ . Let $\phi(x, y, z) := \psi(x, y) \wedge \psi(y, z)$. Since we want the quantifier rank of ϕ to be at most q , we choose n_0 large enough so that $d \log^* n \geq q$ whenever $n \geq n_0$. We claim that this $\phi(x, y, z)$ witnesses the theorem.

Suppose $p(x, y, z)$ is a Boolean combination of quantifier-free formulas and $\text{FO}[\oplus]$ -polynomials of degree bounded by a tower of exponentials of height q/c that agrees with $\phi(x, y, z)$ on more than an ϵ -fraction of graphs with n vertices. Since $q \geq q_0$ is large enough, a tower of exponentials of height $q/c = q/(d+1)$ has height at most $q/d - 4$, and since $q \leq d \log^* n$, the degree is bounded by $\log \log \log \log n$. By Remarks 3.2 and 3.3 we may assume that $p(x, y, z)$ is a Boolean combination of $\text{FO}[\oplus]$ -polynomials of order $2 \log \log \log \log n + 1$, which is at most $\log \log \log n$ if $n \geq n_0$ is large enough.

By Lemma 3.8, we may assume that $p(x, y, z)$ is a regular normal form of this order. If n_0 is large enough, with probability at least $1 - \epsilon/2$, there exists a triple a, b, c of distinct vertices for which $Y(a, b, c)$ holds. Also if n_0 is large enough, with probability at least $1 - \epsilon/2$, the formula $\psi(x, y)$ defines a linear pre-order of width at most 2. By the union bound, with positive probability, all three hold.

- (1) $\phi(x, y, z)$ and $p(x, y, z)$ agree on G .
- (2) $\psi(x, y)$ defines a linear pre-order $\leq$ of width at most 2 in G .
- (3) There exists a triple of distinct vertices a, b, c of G for which $Y(a, b, c)$ holds.

Now assume, without loss of generality, that $a \leq b \leq c$: otherwise, permute them accordingly. Note that we cannot have $c \leq a$, as otherwise, all three a, b, c would belong to the same class of the pre-order, which is not possible, because its width is 2 and a, b, c are distinct. But then $\phi(a, b, c)$ holds and $\phi(c, a, b)$ does not hold, which means that ϕ distinguishes one permutation of (a, b, c) from another. But then p also does; a contradiction to $Y(a, b, c)$. $\square$

Final Remarks. The lower bound is achieved by a formula with free variables. In particular, when we say that $p(x, y, z)$ cannot agree with $\phi(x, y, z)$ on more than an ϵ -fraction of the graphs with n vertices, what we mean is that, on at least a $(1 - \epsilon)$ -fraction of the graphs, the ternary relations on the set of vertices that are defined by $\phi(x, y, z)$ and $p(x, y, z)$ are not identical. It would be nice to obtain a similar kind of lower bound for *sentences*, that is, formulas without free variables. However, since every sentence ϕ agrees on at least half the graphs with n vertices with one of the two constant polynomials $p = 0$ or $p = 1$, the correct lower-bound statement in this setting is different: that for every $\epsilon > 0$ and every sufficiently large n , any low-degree p cannot agree with ϕ on more than a $(\frac{1}{2} + \epsilon)$ -fraction of the graphs with n vertices. A candidate such as sentence ϕ could be the one saying that the number of edges between the minimum and the maximum classes in the pre-order is odd. However, we were not able to prove that this sentence ϕ must be uncorrelated to any low-degree $\text{FO}[\oplus]$ -polynomial p , that is, that for any such p , the probability that ϕ and p agree on a random graph is very close to $1/2$. We leave this as an interesting open problem.

ACKNOWLEDGMENTS

We are grateful to Swastik Kopparty for discussions and comments on a previous version of this article.

REFERENCES

- E. Allender. 1989. A note on the power of threshold circuits. In *Proceedings of the 30th Symposium on Foundations of Computer Science (FOCS)*. 580–584.
- R. Beigel and J. Tarui. 1994. On ACC. *Computational Complex.* 4 (1994), 350–366.
- K. J. Compton. 1989. 0-1 laws in logic and combinatorics. In *NATO Advanced Study Institute on Algorithms and Order*, I. Rival (Ed.), Kluwer, 353–383.

- A. Dawar and E. Grädel. 2010. Properties of almost all graphs and generalized quantifiers. *Fundamenta Informaticae* 98, 4 (2010), 351–372.
- R. Fagin. 1976. Probabilities on finite models. *J. Symbol. Logic* 41, 1 (1976), 50–58.
- Y. V. Glebskiĭ, D. I. Kogan, M. I. Ligon’kiĭ, and V. A. Talanov. 1969. Range and degree of realizability of formulas in the restricted predicate calculus. *Kibernetika* 2 (1969), 17–28.
- L. Hella, P. G. Kolaitis, and K. Luosto. 1996. Almost everywhere equivalence of logics in finite model theory. *Bull. Symbol. Logic* 2 (1996), 422–443.
- R. M. Karp. 1979. Probabilistic analysis of a canonical numbering algorithm for graphs. In *Proceedings of the AMS Symposium in Pure Mathematics*, vol. 34, 365–378.
- Ph. G. Kolaitis and S. Kopparty. 2009a. Random graphs and the parity quantifier. In *Proceedings of the 41st ACM Symposium on the Theory of Computing (STOC)*. 705–714.
- Ph. G. Kolaitis and S. Kopparty. 2009b. Random graphs and the parity quantifier. Technical Report 33, Electronic Colloquium on Computational Complexity (ECCC).
- Ph. G. Kolaitis and M. Y. Vardi. 1992. Infinitary logics and 0-1 laws. *Inform. Computation* 98, 2 (1992), 258–294.
- A. A. Razborov. 1987. Lower bounds on the size of bounded depth networks over a complete basis with logical addition. *Math. Notes Acad. Sci. USSR* 41 (1987), 333–338.
- R. Smolensky. 1987. Algebraic methods in the theory of lower bounds for Boolean circuit complexity. In *Proceedings of the 19th ACM Symposium on the Theory of Computing (STOC)*. 77–82.
- R. Williams. 2011. Non-uniform ACC circuit lower bounds. In *Proceedings of the 26th IEEE Conference on Computational Complexity (CCC)*. 115–125.

Received October 2012; accepted May 2013