Towards Sustainable Forest Management Strategies with MOEAs Appendix A

April 3, 2020

This appendix has the functions and parameter values necessary to use the forest model in [3]. The parameter values are based on the values in [2]. We assume the forest to consist of a single species, which is spruce. The site type is assumed to be Myrtillus (MT), which is the reference type for the model. The site index is $SI = 15$. Thus, the terms and parameter values corresponding to other species or site types are omitted below.

1 The growth model

The trees are divided into $n = 11$ size classes based on the tree diameter d_s . Let γ_s be the basal area per tree in size class s . The values d_s and γ_s corresponding to the size classes are given in Table 1.

The total basal area is computed as

$$B(x_t) = \sum_{s=1}^n \gamma_s x_{st},$$

where x_{st} is the number of trees in size class s at time t within one hectare. The basal area of trees larger than size class s is

$$B_s(x_t) = \sum_{i=s+1}^n \gamma_i x_{it}.$$

The ingrowth function, which is the number of trees that naturally enter the smallest size class, is given by

$$\phi(x_t) = \frac{\exp\left(b_{16} + b_{17}\sqrt{B(x_t)}\right)}{1 + \exp\left(b_{18} - b_{19}\log B(x_t) - b_{22}B(x_t)\right)}, \quad (1)$$

where the parameters b_i are given in Table 2, and $B(x_t)$ is the total basal area of the forest at time t .

Table 1: Size-class specific parameter values for the forest growth model. The table also contains the initial state x_{t_0} .

size class	d_s (cm)	γ_s (m ²)	ν_{1s} (m ³)	ν_{2s} (m ³)	x_{st_0}
1	2.5	0.0005	0	0	0
2	7.5	0.0044	0.0137	0	1750
3	12.5	0.0123	0.0666	0	0
4	17.5	0.0241	0.1669	0	0
5	22.5	0.0398	0.0808	0.2342	0
6	27.5	0.0594	0.0648	0.4458	0
7	32.5	0.0830	0.0598	0.6839	0
8	37.5	0.1106	0.0498	0.9630	0
9	42.5	0.1419	0.0504	1.2531	0
10	47.5	0.1772	0.0432	1.5742	0
11	52.5	0.2165	0.0392	1.8998	0

The fraction of trees in size class s that die during a 5-year period is given by

$$\mu_s(x_t) = 1 - \frac{1}{1 + \exp\left(-b_{10} - b_{11}\sqrt{d_s} - b_{12}d_s - b_{14}\sqrt{B_s(x_t)} - 5b_{15}\right)}.$$

The fraction of trees that move from size class s to the next during a 5-year period is

$$\alpha(x_t) = q^{-1} \exp\left(b_0 + b_1\sqrt{d_s} + b_2d_s + b_3 \log(TS) + b_7 \log B(x_t) + b_9 \frac{B_s(x_t)}{\sqrt{d_s + 1}}\right),$$

where q is the width of a size class (5 cm), and TS is the temperature sum.

2 The Revenue and Cost Functions

Let ν_s be the volume of a tree in size class s (in m³). For the numerical values, we can compute $\nu_s = \nu_{1s} + \nu_{2s}$ using the values in Table 1. The values ν_{1s} and ν_{2s} are the volumes of pulpwood and sawlog in a tree of size class s . The harvesting amounts are denoted by h_{st}

The cost functions for thinning and clearcut are

$$\begin{aligned} C_i(h_t) = & C_{i0}C_{i1} \sum_{s=1}^n h_{st} (C_{i2} + C_{i3}\nu_s - C_{i4}\nu_s^2) \\ & + C_{i5} \left(C_{i6} \sum_{s=1}^n h_{st}\nu_s + C_{i7} \left(\sum_{s=1}^n h_{st}\nu_s \right)^{0.7} \right) \end{aligned}$$

Table 2: Parameter values for the growth model

Parameter	Value
b_0	-9.6448
b_1	0.455
b_2	-0.05741
b_3	1.4551
b_7	-0.3081
b_9	-0.1473
b_{10}	5.871
b_{11}	1.536
b_{12}	-0.122
b_{14}	-0.69
b_{15}	-0.465
b_{16}	4.378
b_{17}	-0.0265
b_{18}	1.001
b_{19}	0.641
b_{22}	-0.0658

Table 3: The parameter values for the cost functions. See [2].

i	C_{i0}	C_{i1}	C_{i2}	C_{i3}	C_{i4}	C_{i5}	C_{i6}	C_{i7}
th	2.1	1.15	0.412	0.758	0.18	1	2.272	0.535
cc	0.1	1	0.397	0.758	0.18	1	1.376	0.393

Table 4: Miscellaneous parameter values for the forest growth model. C_f is the fixed harvesting cost. TS is the temperature sum, representing the climate at the forest site. p_1 and p_2 are the pulpwood and sawlog prices. w is the artificial regeneration cost. r is the interest rate used for discounting.

Parameter	Value
C_f	500 €
TS	1100
p_1	34.07 €
p_2	58.44 €
w	0 €
r	0.03

for $i = th, cc$. For the parameter values, see Table 3. Note, that this differs from the cost function in [3] such that the cutting of the smallest size class is allowed.

The prices for pulpwood and sawlog are p_1 and p_2 , respectively (see Table 4). The revenue is then

$$R(h_t) = \sum_{s=1}^n h_{st}(p_1\nu_{1s} + p_2\nu_{2s}).$$

The objective functions in the article include discounting where the discount factor is

$$b = \frac{1}{1+r},$$

where r is the annual interest rate.

3 Computing the Carbon Storage

The carbon storage is computed using the model as [1]. The computation has the following steps.

The merchantable timber volume of the stand at time period t is

$$\omega_t = \sum_{s=1}^n x_{st}(\nu_{1s} + \nu_{2s})$$

The mass of the merchantable timber is computed using density factor δ . To estimate the mass of the entire tree, we use expansion factor η . The total tree biomass in the stand at time step t is

$$\tilde{B}_t = \rho\eta\omega_t.$$

The dry masses of pulpwood and sawlog harvested at the end of period t are

$$y_{it} = \rho \sum_{s=1}^n h_{st}\nu_{is}$$

for $i = 1, 2$. As trees die naturally, dead tree matter is formed. The dry mass of the dead tree matter formed during a five-year period equals

$$d_{mt} = \rho\eta \sum_{s=1}^n \mu(x_t)_s x_{st}(\nu_{1s} + \nu_{2s}).$$

The dry mass of harvest residues at the end of period t equals

$$d_{ht} = (\eta - 1)(y_{1t} + y_{2t}).$$

The annual decay rates of pulpwood, sawlog, and dead tree matter are denoted by g_1 , g_2 , and g_d , respectively.

Table 5: Parameters used in the calculation of the carbon storage

Parameter	Value
ρ	0.3774
η	2.1566
θ	1.83333
g_1	0.06611
g_2	0.4707
g_d	0.18196

We denote

$$\alpha_i = \frac{g_i}{g_i + r}$$

for $i = 1, 2, m$. This factor represents the future effects of the decaying wood using discounting.

We compute the net negative emissions by the formula

$$Q_t = \theta \left(\tilde{B}_{t+1}(x_{t+1}) - \tilde{B}_t(x_t) + D_1(h_t) + D_2(h_t) + D_d(x_t) \right),$$

where

$$\begin{aligned} D_1(h_t) &= (1 - \alpha_1(r)) y_{1t}(h_t), \\ D_2(h_t) &= (1 - \alpha_2(r)) y_{2t}(h_t), \\ D_d(x_t) &= (1 - \alpha_d(r)) (d_{mt}(x_t) + d_{ht}(h_t)). \end{aligned}$$

The parameter values are in Table 5. This is similar to [1, Equation 6], except that we have not multiplied the net negative emissions by the carbon price.

References

- [1] Aino Assmuth, Janne Rämö, and Olli Tahvonen. Economics of size-structured forestry with carbon storage. *Canadian Journal of Forest Research*, 48(1):11–22, 2017.
- [2] Vesa-Pekka Parkatti, Aino Assmuth, Janne Rämö, and Olli Tahvonen. Economics of boreal conifer species in continuous cover and rotation forestry. *Forest Policy and Economics*, 100:55–67, 2019.
- [3] Timo Pukkala, Erkki Lähde, and Olavi Laiho. Species interactions in the dynamics of even-and uneven-aged boreal forests. *Journal of sustainable forestry*, 32(4):371–403, 2013.